

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 May 2003 (30.05.2003)

PCT

(10) International Publication Number
WO 03/044057 A2

(51) International Patent Classification: C07K 14/705

(21) International Application Number: PCT/IB02/04924

(22) International Filing Date: 29 October 2002 (29.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/332,119 21 November 2001 (21.11.2001) US(71) Applicant (for all designated States except US): GENSET
S.A. [FR/FR]; Intellectual Property Department, 24, rue
Royale, F-75008 Paris (FR).

(72) Inventor; and

(75) Inventor/Applicant (for US only): LUCAS, John
[US/US]; 339 The Valley Road, Concord, MA 01742
(US).(74) Common Representative: GENSET S.A.; c/o De Luca,
Giampiero, Serono International S.A., Intellectual Prop-
erty, 12, Chemin des Aulx, 1228 Plan-les-Ouates (CH).(81) Designated States (national): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: USE OF OBG3 FOR PROMOTING CENTRAL NERVOUS SYSTEM REMYELINATION

(57) Abstract: The present invention relates to the field of central nervous system (CNS) research. Demyelination of neuronal axons within the CNS underlies the pathogenesis of degenerative diseases of the neuromuscular system, such as multiple sclerosis and hereditary leukodystrophies. Therefore, treatments aimed towards accelerating the repair of myelin sheaths offer a potential therapeutic to ameliorate the symptoms of multiple sclerosis and leukodystrophies. A compound, globular OBG3, has been identified that has immunosuppressive properties. This compound should be effective for accelerating the rate of remyelination and treating multiple sclerosis and leukodystrophies.

WO 03/044057 A2

**USE OF OBG3
FOR PROMOTING CENTRAL NERVOUS SYSTEM REMYELINATION**

RELATED APPLICATION INFORMATION

This application claims priority on United States provisional patent application Serial No. 60/332,119, filed November 21, 2000, entitled "Use of OBG3 for promoting central nervous system remyelination".

FIELD OF THE INVENTION

The present invention relates to the field of central nervous system research, in particular the discovery of compounds effective for accelerating the repair of myelin sheaths of demyelinated neurons and useful for treating multiple sclerosis and hereditary leukodystrophies.

BACKGROUND OF THE INVENTION

The following discussion is intended to facilitate the understanding of the invention, but is not intended nor admitted to be prior art to the invention.

Demyelinating diseases are those in which myelin is the primary target. They fall into two main groups: acquired diseases and hereditary metabolic disorders. The most common acquired disease is multiple sclerosis (MS), a chronic, frequently progressive, inflammatory central nervous system (CNS) disease characterized pathologically by primary demyelination, usually without initial axonal injury. The etiology and pathogenesis of MS are unknown. Several immunological features of MS, and its moderate association with certain major histocompatibility complex alleles, have prompted the speculation that MS is an immune-mediated disease.

The hereditary metabolic disorders include eight identified leukodystrophies: metachromatic leukodystrophy, Refsum's disease, adrenoleukodystrophy, Krabbe's disease, phenylketonuria, Canavan disease, Pelizaeus-Merzbacher disease, and Alexander's disease. The first six are storage disorders. The lack of or the malfunctioning of an enzyme causes a toxic buildup of chemical substances. The etiology of Pelizaeus-Merzbacher disease and Alexander's disease, on the other hand, remains unknown.

The limited efficacy of current therapies for MS and other demyelinating diseases has stimulated interest in novel therapies to ameliorate these diseases. However, due to the apparently complex etiopathogenesis of these diseases, potentially involving both environmental and autoimmune factors, the need still exists for an effective treatment of these demyelinating disorders.

SUMMARY OF THE INVENTION

The instant invention is based on the discovery that portions of the full-length OBG3 polypeptide, termed OBG3 polypeptide fragments or gOBG3 polypeptide fragments, have unexpected effects *in vitro* and *in vivo*, including utility for remyelination and immunosuppression in humans and other mammals. These unexpected effects of OBG3 or gOBG3 polypeptide fragment administration in mammals also include reduction of elevated free fatty acid levels caused by administration of epinephrine, *i.v.* injection of "intralipid", or administration of a high fat test meal,

as well as increased fatty acid oxidation in muscle cells, and weight reduction in mammals consuming a high fat/high sucrose diet. These effects are unexpected and surprising given that administration of full-length OBG3 polypeptide typically has no effect or a significantly reduced effect *in vivo* or *in vitro* depending on the specific biological activity and the amount administered.

- 5 To the extent that any effect is observed following administration of full-length OBG3 polypeptide, the levels of full-length OBG3 polypeptide required for an effect render it unfeasible in most instances as a potential treatment for humans at this time. In contrast, the OBG3 and gOBG3 polypeptide fragments of the invention are radically more effective and thus can be provided at levels that are feasible for treatments in humans.

- 10 Thus, the invention is drawn to OBG3 and gOBG3 polypeptide fragments, polynucleotides encoding said OBG3 and gOBG3 polypeptide fragments, vectors comprising said OBG3 and gOBG3 polynucleotides, and cells recombinant for said OBG3 and gOBG3 polynucleotides, as well as to pharmaceutical and physiologically acceptable compositions comprising said OBG3 and gOBG3 polypeptide fragments and methods of administering said OBG3 and gOBG3
15 pharmaceutical and physiologically acceptable compositions in order to accelerate the rate of remyelination in the central nervous system.

- In a first aspect, the invention features a purified, isolated, or recombinant OBG3 or gOBG3 polypeptide fragment that that has significantly greater activity than a full-length OBG3 polypeptide, wherein said activity is accelerating the rate of remyelination of demyelinated neurons in a mammal
20 or human. In preferred embodiments, said polypeptide fragment comprises, consists essentially of, or consists of, at least 6 and not more than 238 consecutive amino acids of SEQ ID NO:6 or at least 6 and not more than 241 consecutive amino acids of SEQ ID NO:2 or SEQ ID NO:4. In other preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 84-244, 85-244, 86-244, 87-244, 88-244, 89-244, 90-244, 91-244, 92-
25 244, 93-244, 94-244, 95-244, 96-244, 97-244, 98-244, 99-244, 100-244, 101-244, 102-244, or 103-244 of SEQ ID NO:6. In other preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 88-247, 89-247, 90-247, 91-247, 92-247, 93-247, 94-247, 95-247, 96-247, 97-247, 98-247, 99-247, 100-247, 101-247, 102-247, 103-247, 104-247, 105-247, or 106-247 of SEQ ID NO:2 or SEQ ID NO:4. In other preferred embodiments,
30 OBG3 or gOBG3 polypeptide fragments are selected from amino acids about 84 to 244, 85 to 244, 101 to 244, 102 to 244, or 103 to 244 of SEQ ID NO:6 and amino acids 88 to 247, 104 to 247, 105 to 247, or 106 to 247 of SEQ ID NO:2 or SEQ ID NO:4. In further preferred embodiments, gOBG3 polypeptide fragments are said selected gOBG3 polypeptide fragments made resistant to dipeptidyl peptidase cleavage by N-terminal modification. In other further preferred embodiments, said
35 polypeptide fragment comprises an amino acid sequence at least 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the corresponding consecutive amino acids of SEQ ID NO:6, SEQ ID NO:2 or SEQ ID NO:4.

In other highly preferred embodiments, said polypeptide fragment comprises, consists essentially of, or consists of, a purified, isolated, or recombinant gOBG3 fragment. Preferably, said gOBG3 polypeptide fragment comprises, consists essentially of, or consists of, at least 6 consecutive amino acids of amino acids 84 to 244 of SEQ ID NO:6 or at least 6 consecutive amino acids of amino acids 88 to 247 of SEQ ID NO:2 or SEQ ID NO:4. In other preferred embodiments, gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 84-244, 85-244, 86-244, 87-244, 88-244, 89-244, 90-244, 91-244, 92-244, 93-244, 94-244, 95-244, 96-244, 97-244, 98-244, 99-244, 100-244, 101-244, 102-244, or 103-244 of SEQ ID NO:6. In other preferred embodiments, gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 88-247, 89-247, 90-247, 91-247, 92-247, 93-247, 94-247, 95-247, 96-247, 97-247, 98-247, 99-247, 100-247, 101-247, 102-247, 103-247, 104-247, 105-247, or 106-247 of SEQ ID NO:2 or SEQ ID NO:4. In other preferred embodiments, gOBG3 polypeptide fragments are selected from amino acids 84 to 244, 85 to 244, 101 to 244, 102 to 244, or 103 to 244 of SEQ ID NO:6 and amino acids 88 to 247, 104 to 247, 105 to 247, or 106 to 247 of SEQ ID NO:2 or SEQ ID NO:4. In further preferred embodiments, gOBG3 polypeptide fragments are said selected gOBG3 polypeptide fragments made resistant to dipeptidyl peptidase cleavage by N-terminal modification. Alternatively, said gOBG3 fragment comprises, consists essentially of, or consists of, an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the corresponding amino acids 84 to 244 of SEQ ID NO:6 or at least 75% identical to amino acids 88 to 247 of SEQ ID NO:2 or SEQ ID NO:4. In a further preferred embodiment, the OBG3 polypeptide fragment is identical to an APM1 proteolytic cleavage product from human plasma. Preferably, the proteolytic cleavage product comprises the C1q globular head or a portion thereof. More preferably, the proteolytic cleavage product is identical to a proteolytic cleavage product isolated from human plasma by immunoprecipitation using antibodies specific for the C1q globular head. More preferably the proteolytic cleavage product cannot be immunoprecipitated from human plasma using an antibody directed against the human non-homologous region (HDQETTTQGGVLLPLPKGA) of APM1. Still more preferably, the APM1 proteolytic cleavage product has an apparent molecular weight of 27 kDa using SDS-PAGE.

In another aspect, the present invention is also directed towards polypeptide fragments of OBG3 having biological activity. Preferred polypeptide fragment is amino acids 166-193 or further fragments thereof. Particularly preferred polypeptide fragment is amino acids 166-176. Other particularly preferred fragment is 167-176.

In a further preferred embodiment, the OBG3 or gOBG3 polypeptide fragment is able to increase the rate of cellular division of oligodendrocyte progenitor cells in a mammal or human.

Further preferred OBG3 or gOBG3 polypeptide fragments are those that increase the rate of migration of oligodendrocyte progenitor cells from the germinal centers to demyelinated axons in the CNS of a mammal or human.

Further preferred OBG3 or gOBG3 polypeptide fragments are those that induce oligodendrocyte progenitor cells to differentiate into mature dendrocytic cells at sites of demyelination in axons of in the CNS of a mammal or a human.

Further preferred OBG3 or gOBG3 polypeptide fragments are those that significantly reduce or eliminate the symptoms of multiple sclerosis in a mammal or human.

Further preferred OBG3 or gOBG3 polypeptide fragments are those that significantly reduce or eliminate the symptoms of hereditary leukodystrophies, which include metachromatic leukodystrophy, Refsum's disease, adrenoleukodystrophy, Krabbe's disease, phenylketonuria, Canavan disease, Pelizaeus-Merzbacher disease, and Alexander's disease.

Further preferred OBG3 or gOBG3 polypeptide fragments are those that form multimers (e.g., heteromultimers or homomultimers) *in vitro* and/or *in vivo*. Preferred multimers are homodimers or homotrimer. Other preferred multimers are homomultimers comprising at least 4, 6, 8, 9, 10, or 12 OBG3 or gOBG3 polypeptide fragment subunits. Other preferred mulimers are hetero multimers comprising a OBG3 or gOBG3 polypeptide fragment of the invention.

Further preferred embodiments include heterologous polypeptides comprising an OBG3 or gOBG3 polypeptide fragment of the invention.

In a further embodiment, the invention features a pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of, said OBG3 or gOBG3 polypeptide fragment described in the first aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

In a further embodiment, the invention features a method for the production of a pharmaceutical or physiologically acceptable composition consisting essentially of, or consisting of, said OBG3 or gOBG3 polypeptide fragment described in the first aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

Full-length OBG3 (ACRP30, AdipoQ, APM1) polypeptides and polynucleotides encoding the same may be specifically substituted for an OBG3 or gOBG3 polypeptide fragment or polynucleotide encoding the same in any embodiment of the present invention.

PREFERRED EMBODIMENTS OF THE INVENTION

I. OBG3 Polypeptide Fragments of the Invention

The invention is drawn, *inter alia*, to isolated, purified or recombinant OBG3 polypeptide fragments. OBG3 polypeptide fragments of the invention are useful for accelerating the regeneration of myelin sheaths *in vivo*. OBG3 polypeptide fragments are also useful *inter alia* in screening assays for agonists or antagonists of OBG3 fragment activity; in screening assays for antagonists of dipeptidyl peptidase cleavage of OBG3 fragments, preferably cleavage of the N-

terminal EP dipeptide of OBG3 polypeptide fragment 103-244 of SEQ ID NO:6, cleavage of the N-terminal VP dipeptide of OBG3 fragment 85-244 of SEQ ID NO:6, or cleavage of the N-terminal EP dipeptide of OBG3 polypeptide fragment 106-247 of SEQ ID NO:2 or SEQ ID NO:4; for raising OBG3 fragment-specific antibodies; and in diagnostic assays.

5 The full-length OBG3 polypeptide is comprised of at least four distinct regions including:

1. an N-terminal putative signal sequence about from amino acids 1-17 of SEQ ID NO:6, SEQ ID NO:2, or SEQ ID NO:4;
2. a unique region about from amino acids 18-41 of SEQ ID NO:6 or 18-44 of SEQ ID NO:2, or SEQ ID NO:4;
- 10 3. a collagen-like region about from amino acids 42-107 of SEQ ID NO:6 or 45-110 of SEQ ID NO:2 or SEQ ID NO:4; and
4. a globular region about from amino acids 108-244 of SEQ ID NO:6 or 111-247 of SEQ ID NO:2 or SEQ ID NO:4.

15 The term "collagen residues" is used in the manner standard in the art to mean the amino acid triplet glycine, X, Y, where X and Y can be any amino acid.

The OBG3 polypeptide fragments of the present invention are preferably provided in an isolated form, and may be partially or substantially purified. A recombinantly produced version of an OBG3 polypeptide fragment can be substantially purified by the one-step method described by Smith et al. ((1988) Gene 67(1):31-40) or by the methods described herein or known in the art (see, 20 e.g., Examples 1-3). Fragments of the invention also can be purified from natural or recombinant sources using antibodies directed against the polypeptide fragments of the invention by methods known in the art of protein purification.

Preparations of OBG3 polypeptide fragments of the invention involving a partial purification of or selection for the OBG3 polypeptide fragments are also specifically contemplated. 25 These crude preparations are envisioned to be the result of the concentration of cells expressing OBG3 polypeptide fragments with perhaps a few additional purification steps, but prior to complete purification of the fragment. The cells expressing OBG3 polypeptide fragments are present in a pellet, they are lysed, or the crude polypeptide is lyophilized, for example.

OBG3 or gOBG3 polypeptide fragments can be any integer in length from at least 6 30 consecutive amino acids to 1 amino acid less than a full-length OBG3 polypeptide. Thus, for human OBG3 (SEQ ID NO: 6), an OBG3 or gOBG3 polypeptide fragment can be any integer of consecutive amino acids from 6 to 243; for mouse OBG3 (SEQ ID NO:2 or SEQ ID NO:4) an OBG3 or gOBG3 fragment can be any integer of consecutive amino acids from 6 to 246, for example. The term "integer" is used herein in its mathematical sense and thus representative 35 integers include: 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82,

83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, and 246.

Each OBG3 fragment as described above can be further specified in terms of its N-terminal and C-terminal positions. For example, every combination of a N-terminal and C-terminal position that fragments of from 6 contiguous amino acids to 1 amino acids less than the full-length OBG3 polypeptide could occupy, on any given intact and contiguous full-length OBG3 polypeptide sequence are included in the present invention. Thus, a 6 consecutive amino acid fragment could occupy positions selected from the group consisting of 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 7-12, 8-13, 9-14, 10-15, 11-16, 12-17, 13-18, 14-19, 15-20, 16-21, 17-22, 18-23, 19-24, 20-25, 21-26, 22-27, 23-28, 24-29, 25-30, 26-31, 27-32, 28-33, 29-34, 30-35, 31-36, 32-37, 33-38, 34-39, 35-40, 36-41, 37-42, 38-43, 39-44, 40-45, 41-46, 42-47, 43-48, 44-49, 45-50, 46-51, 47-52, 48-53, 49-54, 50-55, 51-56, 52-57, 53-58, 54-59, 55-60, 56-61, 57-62, 58-63, 59-64, 60-65, 61-66, 62-67, 63-68, 64-69, 65-70, 66-71, 67-72, 68-73, 69-74, 70-75, 71-76, 72-77, 73-78, 74-79, 75-80, 76-81, 77-82, 78-83, 79-84, 80-85, 81-86, 82-87, 83-88, 84-89, 85-90, 86-91, 87-92, 88-93, 89-94, 90-95, 91-96, 92-97, 93-98, 94-99, 95-100, 96-101, 97-102, 98-103, 99-104, 100-105, 101-106, 102-107, 103-108, 104-109, 105-110, 106-111, 107-112, 108-113, 109-114, 110-115, 111-116, 112-117, 113-118, 114-119, 115-120, 116-121, 117-122, 118-123, 119-124, 120-125, 121-126, 122-127, 123-128, 124-129, 125-130, 126-131, 127-132, 128-133, 129-134, 130-135, 131-136, 132-137, 133-138, 134-139, 135-140, 136-141, 137-142, 138-143, 139-144, 140-145, 141-146, 142-147, 143-148, 144-149, 145-150, 146-151, 147-152, 148-153, 149-154, 150-155, 151-156, 152-157, 153-158, 154-159, 155-160, 156-161, 157-162, 158-163, 159-164, 160-165, 161-166, 162-167, 163-168, 164-169, 165-170, 166-171, 167-172, 168-173, 169-174, 170-175, 171-176, 172-177, 173-178, 174-179, 175-180, 176-181, 177-182, 178-183, 179-184, 180-185, 181-186, 182-187, 183-188, 184-189, 185-190, 186-191, 187-192, 188-193, 189-194, 190-195, 191-196, 192-197, 193-198, 194-199, 195-200, 196-201, 197-202, 198-203, 199-204, 200-205, 201-206, 202-207, 203-208, 204-209, 205-210, 206-211, 207-212, 208-213, 209-214, 210-215, 211-216, 212-217, 213-218, 214-219, 215-220, 216-221, 217-222, 218-223, 219-224, 220-225, 221-226, 222-227, 223-228, 224-229, 225-230, 226-231, 227-232, 228-233, 229-234, 230-235, 231-236, 232-237, 233-238, 234-239, 235-240, 236-241, 237-242, 238-243, and 239-244 of SEQ ID NO:6. A 238 consecutive amino acid fragment could occupy positions selected from the group consisting of 1-238, 2-239, 3-240, 4-241, 5-242, 6-243 and 7-244 of SEQ ID NO:6. Similarly, the positions occupied by all the other fragments of sizes between 6 amino acids and 243 amino acids on

SEQ ID NO:6 are included in the present invention and can also be immediately envisaged based on these two examples and therefore, are not individually listed solely for the purpose of not unnecessarily lengthening the specification. Furthermore, the positions occupied by fragments of 6 to 241 consecutive amino acids on SEQ ID NO:2 or SEQ ID NO:4 are included in the present invention and can also be immediately envisaged based on these two examples and therefore are not individually listed solely for the purpose of not unnecessarily lengthening the specification. In addition, the positions occupied by fragments of 6 consecutive amino acids to 1 amino acid less than any other full-length OBG3 polypeptide can also be envisaged based on these two examples and therefore are not individually listed solely for the purpose of not unnecessarily lengthening the specification. In preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 84-244, 85-244, 86-244, 87-244, 88-244, 89-244, 90-244, 91-244, 92-244, 93-244, 94-244, 95-244, 96-244, 97-244, 98-244, 99-244, 100-244, 101-244, 102-244, or 103-244 of SEQ ID NO:6. In preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 88-247, 89-247, 90-247, 91-247, 92-247, 93-247, 94-247, 95-247, 96-247, 97-247, 98-247, 99-247, 100-247, 101-247, 102-247, 103-247, 104-247, 105-247, or 106-247 of SEQ ID NO:2 or SEQ ID NO:4.

The OBG3 or gOBG3 polypeptide fragments of the present invention may alternatively be described by the formula "n to c" (inclusive); where "n" equals the N-terminal most amino acid position (as defined by the sequence listing) and "c" equals the C-terminal most amino acid position (as defined by the sequence listing) of the polypeptide; and further where "n" equals an integer between 1 and the number of amino acids of the full length polypeptide sequence of the present invention minus 6 (238 for SEQ ID NO: 6 and 241 for SEQ ID NOs: 2 or 4); and where "c" equals an integer between 7 and the number of amino acids of the full-length polypeptide sequence (244 for SEQ ID NO: 6 and 247 for SEQ ID NOs: 2 or 4); and where "n" is an integer smaller than "c" by at least 6. Therefore, for SEQ ID NO: 6, "n" is any integer selected from the list consisting of: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 234, 235, 236, 237 and 238; and "c" is any integer selected from the group consisting of: 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33,

34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244. Every combination of "n" and "c" positions are included as specific embodiments of the invention. Moreover, the formula "n" to "c" may be modified as "n1 - n2" to "c1 - c2", wherein "n1 - n2" and "c1 - c2" represent positional ranges selected from any two integers above which represent amino acid positions of the sequence listing. Alternative formulas include "n1 - n2" to "c" and "n" to "c1 - c2". In preferred embodiment, OBG3 or gOBG3 polypeptide fragments of the invention may be described by the formula where n1=84, n2=103, and c=244 of SEQ ID NO:6 or by the formula n1=88, n2=106, and c=247 of SEQ ID NO: 2 or SEQ ID NO:4.

These specific embodiments, and other polypeptide and polynucleotide fragment embodiments described herein may be modified as being "at least", "equal to", "equal to or less than", "less than", "at least ___ but not greater than ___" or "from ___ to ___", a specified size or specified N-terminal and/or C-terminal positions. It is noted that all ranges used to describe any embodiment of the present invention are inclusive unless specifically set forth otherwise.

The present invention also provides for the exclusion of any individual fragment specified by N-terminal and C-terminal positions or of any fragment specified by size in amino acid residues as described above. In addition, any number of fragments specified by N-terminal and C-terminal positions or by size in amino acid residues as described above may be excluded as individual species. Further, any number of fragments specified by N-terminal and C-terminal positions or by size in amino acid residues as described above may make up a polypeptide fragment in any combination and may optionally include non-OBG3 polypeptide sequence as well.

In particularly preferred embodiments, the OBG3 polypeptide fragment is a "globular OBG3" (gOBG3) fragment. The term "gOBG3 fragment" or "gOBG3" or "gOBG3 polypeptide" as used herein refers to fragments of a full-length OBG3 polypeptide that comprise at least 6 and any other integer number of amino acids up to 137 of the globular region of a full-length OBG3 polypeptide (defined above). In preferred embodiments, gOBG3 polypeptide fragments also comprise at least 1 and any other integer number of amino acids up to 66 of the collagen region of a full-length OBG3 polypeptide, preferably 1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22, 23, or 24 consecutive amino acid residues from the collagen region of the intact OBG3 polypeptide

that are adjacent to the globular region. By "adjacent" to the globular region is meant the first collagen amino acid immediately N-terminal to the globular region and adding each collagen amino acid consecutively in the N-terminal direction. Thus, for example, if there is only one collagen amino acid in the gOBG3 polypeptide fragment, it is the collagen amino acid 107 of SEQ ID NO: 6 or amino acid 110 of SEQ ID NO:2 or SEQ ID NO:4 located adjacent and 5' to the first amino acid of the globular region. If there are 24 collagen amino acids adjacent to the globular region in the gOBG3 fragment they would be the collagen amino acids 84-107 of SEQ ID NO: 6 or amino acids 87-110 of SEQ ID NO:2 or SEQ ID NO:4.

In preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 84-244, 85-244, 86-244, 87-244, 88-244, 89-244, 90-244, 91-244, 92-244, 93-244, 94-244, 95-244, 96-244, 97-244, 98-244, 99-244, 100-244, 101-244, 102-244, or 103-244 of SEQ ID NO:6. In other preferred embodiments, OBG3 or gOBG3 polypeptide fragments having unexpected activity are selected from amino acids 88-247, 89-247, 90-247, 91-247, 92-247, 93-247, 94-247, 95-247, 96-247, 97-247, 98-247, 99-247, 100-247, 101-247, 102-247, 103-247, 104-247, 105-247, or 106-247 of SEQ ID NO:2 or SEQ ID NO:4. In further preferred embodiments, gOBG3 polypeptide fragments are selected from amino acids 84 to 244, 85 to 244, 101 to 244, 102 to 244, or 103 to 244 of SEQ ID NO:6 and amino acids 88 to 247, 104 to 247, 105 to 247, or 106 to 247 of SEQ ID NO:2 or SEQ ID NO:4. In yet other preferred embodiments, the invention features a gOBG3 polypeptide fragment comprising at least 115, but not more than 175 contiguous amino acids of any one of the gOBG3 fragment sequences set forth in Figure 1, wherein no more than 24 of said at least 115 and no more than 175 contiguous amino acids are present in the collagen-like region of OBG3. Preferably, the gOBG3 polypeptide fragment comprises at least 125, but not more than 165, or at least 140, but not more than 165 amino acids, and no more than 24 amino acids are in the collagen-like region; more preferably at least 125 but not more than 165, or at least 140 but not more than 165 amino acids, and no more than 12 amino acids are in the collagen-like region; or at least 140 and not more than 150 amino acids, and no more than 8 amino acids are present in the collagen-like region. Preferably the gOBG3 fragment is mammalian, preferably human or mouse, but most preferably human.

OBG3 and gOBG3 polypeptide fragments of the invention include variants, fragments, analogs and derivatives of the OBG3 and gOBG3 polypeptide fragments described above, including modified OBG3 and gOBG3 polypeptide fragments. Particularly preferred are proteolytically cleaved fragments of OBG3 of SEQ ID NO:6, SEQ ID NO:2, or SEQ ID NO:4. More preferred is OBG3 fragment of about amino acids 85-244 of SEQ ID NO:6 made by collagenase cleavage of SEQ ID NO:6 at about position 84. More preferred is OBG3 fragment of about amino acids 88-247 of SEQ ID NO:2 or of SEQ ID NO:4 made by collagenase cleavage of SEQ ID NO:2 or SEQ ID NO:4 at about position 87. More preferred is OBG3 fragment of about amino acids 85-244 of SEQ ID NO:6 made by matrix metalloproteinase-1 (MMP-1) cleavage of SEQ ID NO:6 at about position

84. More preferred is OBG3 fragment of about amino acids 88-247 of SEQ ID NO:2 or of SEQ ID NO:4 made by matrix metalloproteinase-1 (MMP-1) cleavage of SEQ ID NO:2 or SEQ ID NO:4 at about position 87. More preferred is OBG3 fragment of about amino acids 101-244 of SEQ ID NO:6 made by plasmin cleavage of SEQ ID NO:6 at about position 100. More preferred is OBG3
5 fragment of about amino acids 104-247 of SEQ ID NO:2 or of SEQ ID NO:4 made by plasmin cleavage of SEQ ID NO:2 or SEQ ID NO:4 at about position 103. More preferred is OBG3 fragment of about amino acids 103-244 of SEQ ID NO:6 made by precerebellin processing protease cleavage of SEQ ID NO:6 at about position 102. More preferred is OBG3 fragment of about amino acids 166-247 of SEQ ID NO:2 or of SEQ ID NO:4 made by precerebellin processing protease
10 cleavage of SEQ ID NO:2 or SEQ ID NO:4 at about position 105. More preferred is APM1 proteolytic fragment of SEQ ID NO:6, wherein said APM1 fragment isolated from human plasma migrates with an apparent molecular weight of about 27 kDa on SDS-PAGE under reducing conditions. Most preferred are polypeptide fragments possessing biological activity, comprising amino acids 166-193, amino acids 166-176, and amino acids 167-176

15 Variants

It will be recognized by one of ordinary skill in the art that some amino acids of the OBG3 and gOBG3 fragment sequences of the present invention can be varied without significant effect on the structure or function of the protein; there will be critical amino acids in the fragment sequence that determine activity. Thus, the invention further includes variants of OBG3 and gOBG3
20 polypeptide fragments that have remyelinating activity as described above. Such variants include OBG3 fragment sequences with one or more amino acid deletions, insertions, inversions, repeats, and substitutions either from natural mutations or human manipulation selected according to general rules known in the art so as to have little effect on activity. Guidance concerning how to make phenotypically silent amino acid substitutions is provided below.

25 There are two main approaches for studying the tolerance of an amino acid sequence to change (*see*, Bowie, et al. (1990) Science, 247, 1306-10). The first method relies on the process of evolution, in which mutations are either accepted or rejected by natural selection. The second approach uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene and selections or screens to identify sequences that maintain functionality.

30 These studies have revealed that proteins are surprisingly tolerant of amino acid substitutions and indicate which amino acid changes are likely to be permissive at a certain position of the protein. For example, most buried amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Other such phenotypically silent substitutions are described by Bowie et al. (*supra*) and the references cited therein.

35 Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Phe; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln;

exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe, Tyr. In addition, the following groups of amino acids generally represent equivalent changes: (1) Ala, Pro, Gly, Glu, Asp, Gln, Asn, Ser, Thr; (2) Cys, Ser, Tyr, Thr; (3) Val, Ile, Leu, Met, Ala, Phe; (4) Lys, Arg, His; (5) Phe, Tyr, Trp, His.

5 Similarly, amino acids in the OBG3 and gOBG3 polypeptide fragment sequences of the invention that are essential for function can also be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (*see, e.g.*, Cunningham, et al. (1989) Science 244(4908):1081-5). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for obesity-related activity
10 using assays as described above. Of special interest are substitutions of charged amino acids with other charged or neutral amino acids that may produce proteins with highly desirable improved characteristics, such as less aggregation. Aggregation may not only reduce activity but also be problematic when preparing pharmaceutical or physiologically acceptable formulations, because aggregates can be immunogenic (*see, e.g.*, Pinckard, et al., (1967) Clin. Exp. Immunol 2:331-340;
15 Robbins, et al., (1987) Diabetes Jul;36(7):838-41; and Cleland, et al., (1993) Crit Rev Ther Drug Carrier Syst. 10(4):307-77).

Thus, the fragment, derivative, analog, or homolog of the OBG3 or gOBG3 fragment of the present invention may be, for example: (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino
20 acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code (i.e. may be a non-naturally occurring amino acid); or (ii) one in which one or more of the amino acid residues includes a substituent group; or (iii) one in which the OBG3 or gOBG3 fragment is fused with another compound, such as a compound to increase the half-life of the fragment (for example, polyethylene glycol); or (iv) one in which the additional amino acids are fused to the above
25 form of the fragment, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the above form of the fragment or a pro-protein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

A further embodiment of the invention relates to a polypeptide which comprises the amino
30 acid sequence of an OBG3 or gOBG3 polypeptide fragment having an amino acid sequence which contains at least one conservative amino acid substitution, but not more than 50 conservative amino acid substitutions, not more than 40 conservative amino acid substitutions, not more than 30 conservative amino acid substitutions, and not more than 20 conservative amino acid substitutions. Also provided are polypeptides which comprise the amino acid sequence of a OBG3 or gOBG3
35 fragment, having at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 conservative amino acid substitutions.

Another specific embodiment of a modified OBG3 or gOBG3 fragment of the invention is a polypeptide that is resistant to proteolysis, for example a OBG3 or gOBG3 fragment in which a -CONH- peptide bond is modified and replaced by one or more of the following: a (CH₂NH) reduced bond; a (NHCO) retro inverse bond; a (CH₂-O) methylene-oxy bond; a (CH₂-S) thiomethylene bond; a (CH₂CH₂) carba bond; a (CO-CH₂) cetomethylene bond; a (CHOH-CH₂) hydroxyethylene bond; a (N-N) bound; a E-alcene bond; or a -CH=CH- bond. Thus, the invention also encompasses an OBG3 or gOBG3 fragment or a variant thereof in which at least one peptide bond has been modified as described above.

A further embodiment of the invention relates to an OBG3 or gOBG3 polypeptide fragment made resistant to dipeptidyl peptidase cleavage through N-terminal modification of said polypeptide fragment. In preferred embodiment, said OBG3 or gOBG3 polypeptide fragment is selected from amino acids 85-244 or 103-244 of SEQ ID NO:6 or amino acids 106-247 of SEQ ID NO:2 or SEQ ID NO:4. In preferred embodiment, said dipeptidyl peptidase cleavage leads to removal of the N-terminal dipeptide EP by dipeptidyl peptidase from said preferred gOBG3 polypeptide fragment 103-244 of SEQ ID NO:6 or 106-247 of SEQ ID NO:2 or SEQ ID NO:4. In preferred embodiment, said dipeptidyl peptidase cleavage leads to removal of the N-terminal dipeptide VP by dipeptidyl peptidase from said preferred gOBG3 polypeptide fragment 85-244 of SEQ ID NO:6. In preferred embodiment, said dipeptidyl peptidase is human plasma comprised of dipeptidyl peptidase. In preferred embodiment, said dipeptidyl peptidase is selected from but not restricted to human CD26 or human Attractin. In further preferred embodiment, said dipeptidyl peptidase is selected from soluble human CD26 or soluble human Attractin. In preferred embodiment, said N-terminal modification is selected from but not restricted to glycation [Harte (2001) Regulatory Peptides 96:95-104 which disclosure is hereby incorporated by reference in its entirety], N-methylation, alpha-methylation, desamidation [Gallwitz (2000) Regulatory Peptides 86:103-111 which disclosure is hereby incorporated by reference in its entirety], or alternation of the chirality of one or more N-terminal amino acids [Siegel (1999) European Journal of Clinical Investigation 29:610-614 which disclosure is hereby incorporated by reference in its entirety]. Thus, the invention also encompasses an OBG3 or gOBG3 polypeptide fragment or a variant thereof that has been made resistant to dipeptidyl peptidase cleavage through N-terminal modification of said polypeptide fragment.

In addition, amino acids have chirality within the body of either L or D. In some embodiments it is preferable to alter the chirality of the amino acids in the OBG3 or gOBG3 polypeptide fragments of the invention in order to extend half-life within the body. In other embodiments, it is preferable to alter the chirality of one or more amino acid in order to render the OBG3 or gOBG3 polypeptide fragment resistant to dipeptidyl peptidase cleavage [Siegel (1999) European Journal of Clinical Investigation 29:610-614 which disclosure is hereby incorporated by reference in its entirety]. In further embodiments, it is preferable to alter the chirality of the penultimate N-terminal amino acid in order to render the OBG3 or gOBG3 polypeptide fragment

resistant to dipeptidyl peptidase cleavage. Thus, in some embodiments, one or more of the amino acids are preferably in the L configuration. In other embodiments, one or more of the amino acids are preferably in the D configuration.

Percent Identity

5 The polypeptides of the present invention also include polypeptides having an amino acid sequence at least 50% identical, at least 60% identical, or 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an OBG3 or gOBG3 fragment as described above. By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to an OBG3 or gOBG3 fragment amino acid sequence is meant that the amino acid sequence is identical to the
10 OBG3 or gOBG3 polypeptide fragment sequence except that it may include up to five amino acid alterations per each 100 amino acids of the OBG3 or gOBG3 polypeptide fragment amino acid sequence. The reference sequence is the OBG3 or gOBG3 polypeptide fragment with a sequence corresponding to the sequence of the sequence listing. Thus, to obtain a polypeptide having an amino acid sequence at least 95% identical to an OBG3 or gOBG3 fragment amino acid sequence,
15 up to 5% (5 of 100) of the amino acid residues in the sequence may be inserted, deleted, or substituted with another amino acid compared with the OBG3 or gOBG3 polypeptide fragment sequence. These alterations may occur at the amino or carboxy termini or anywhere between those terminal positions, interspersed either individually among residues in the sequence or in one or more contiguous groups within the sequence.

20 As a practical matter, whether any particular polypeptide is a percentage identical to an OBG3 or gOBG3 fragment can be determined conventionally using known computer programs. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, (1988) Proc Natl Acad Sci USA 85(8):2444-8; Altschul et al., (1990) J Mol Biol 215(3):403-410; Thompson et al., (1994) Nucleic
25 Acids Res 22(2):4673-4680; Higgins et al., (1996) Meth Enzymol 266:383-402; Altschul et al., (1997) Nuc Acids Res 25:3389-3402; Altschul et al., (1993) Nature Genetics 3:266-272). In a particularly preferred embodiment, protein and nucleic acid sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST"), which is well known in the art (*See, e.g.*, Karlin and Altschul (1990) Proc Natl Acad Sci USA 87(6):2264-8; Altschul et al., 1990, 1993, 1997,
30 all supra). In particular, five specific BLAST programs are used to perform the following tasks:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;
- 35 (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

(4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and

(5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

5 The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (*i.e.*, aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62
10 matrix (see, Gonnet et al., (1992) Science 256(5062):1443-5; Henikoff and Henikoff (1993) Proteins 17(1):49-61). Less preferably, the PAM or PAM250 matrices may also be used (See, *e.g.*, Schwartz and Dayhoff, eds, (1978) Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure, Washington: National Biomedical Research Foundation). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably
15 selects those segments which satisfy a user-specified threshold of significance, such as a user-specified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (See, *e.g.*, Karlin and Altschul, (1990) Proc Natl Acad Sci USA 87(6):2264-8). The BLAST programs may be used with the default parameters or with modified parameters provided by the user. Preferably, the parameters are default
20 parameters.

A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (1990) Comp. App. Biosci. 6:237-245. In a sequence alignment the query and subject
25 sequences are both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group=25 Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=247 or the length of the subject amino acid sequence, whichever is shorter.

30 If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, the results, in percent identity, must be manually corrected because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues
35 of the query sequence that are N- and C- terminal of the subject sequence, that are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence

alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query amino acid residues outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100-residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not match/align with the first residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%.

In another example, a 90-residue subject sequence is compared with a 100-residue query sequence. This time the deletions are internal so there are no residues at the N- or C-termini of the subject sequence, which are not matched/aligned with the query. In this case, the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected. No other manual corrections are made for the purposes of the present invention.

Production

Note, throughout the disclosure, wherever OBG3 polypeptide fragments are discussed, gOBG3 fragments are specifically intended to be included as a preferred subset of OBG3 polypeptide fragments.

OBG3 polypeptide fragments are preferably isolated from human or mammalian tissue samples or expressed from human or mammalian genes in human or mammalian cells. The OBG3 polypeptide fragments of the invention can be made using routine expression methods known in the art. The polynucleotide encoding the desired polypeptide fragments is ligated into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems are used in forming recombinant polypeptide fragments. The polypeptide fragment is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Purification is by any technique known in the art, for example, differential extraction, salt fractionation, chromatography, centrifugation, and the like. See, for example, Methods in Enzymology for a variety of methods for purifying proteins. Also, see Examples 1-3 for methods previously used for OBG3 polypeptide fragments.

In an alternative embodiment, the polypeptides of the invention are isolated from milk. The polypeptides can be purified as full-length OBG3 polypeptides, which can then be cleaved, if appropriate, in vitro to generate an OBG3 fragment, or, alternatively, OBG3 fragments themselves can be purified from the milk. Any of a large number of methods can be used to purify the present polypeptides from milk, including those taught in Protein Purification Applications, A Practical Approach (New Edition), Edited by Simon Roe, AEA Technology Products and Systems, Biosciences, Harwell; Clark (1998) J Mammary Gland Biol Neoplasia 3:337-50; Wilkins and Velander (1992) 49:333-8; U.S. Patent Nos. 6,140,552; 6,025,540; Hennighausen, Protein Expression and Purification, vol. 1, pp. 3-8 (1990); Harris et al. (1997) Bioseparation 7:31-7; Degener et al. (1998) J Chromatog 799:125-37; Wilkins (1993) J Cell Biochem Suppl. 0 (17 part A):39; the entire disclosures of each of which are herein incorporated by reference. In a typical embodiment, milk is centrifuged, e.g. at a relatively low speed, to separate the lipid fraction, and the aqueous supernatant is then centrifuged at a higher speed to separate the casein in the milk from the remaining, "whey" fraction. Often, biomedical proteins are found in this whey fraction, and can be isolated from this fraction using standard chromatographic or other procedures commonly used for protein purification, e.g. as described elsewhere in the present application. In one preferred embodiment, OBG3 polypeptides are purified using antibodies specific to OBG3 polypeptides, e.g. using affinity chromatography. In addition, methods can be used to isolate particular OBG3 fragments, e.g. electrophoretic or other methods for isolating proteins of a particular size. The OBG3 polypeptides isolating using these methods can be naturally occurring, as OBG3 polypeptides have been discovered to be naturally present in the milk of mammals (see, e.g. Example 17), or can be the result of the recombinant production of the protein in the mammary glands of a non-human mammal, as described infra. In one such embodiment, the OBG3 fragment is produced as a fusion protein with a heterologous, antigenic polypeptide sequence, which antigenic sequence can be used to purify the protein, e.g., using standard immuno-affinity methodology.

In addition, shorter protein fragments may be produced by chemical synthesis. Alternatively, the proteins of the invention are extracted from cells or tissues of humans or non-human animals. Methods for purifying proteins are known in the art, and include the use of detergents or chaotropic agents to disrupt particles followed by differential extraction and separation of the polypeptides by ion exchange chromatography, affinity chromatography, sedimentation according to density, and gel electrophoresis.

Any OBG3 fragment cDNA, including that in Fig. 4, can be used to express OBG3 polypeptide fragments. The nucleic acid encoding the OBG3 fragment to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The OBG3 fragment cDNA insert in the expression vector may comprise the coding sequence for: the full-length OBG3 polypeptide (to be later modified); from 6 amino acids to 6 amino acids less than the full-length OBG3 polypeptide; a gOBG3 fragment; or variants and % similar polypeptides.

The expression vector is any of the mammalian, yeast, insect or bacterial expression systems known in the art, some of which are described herein, and examples of which are given in the Examples (Examples 1-3). Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and facilitate proper protein folding, the codon context and codon pairing of the sequence can be optimized for the particular expression organism into which the expression vector is introduced, as explained by Hatfield, et al., US Patent Number 5,082,767, the disclosures of which are incorporated by reference herein in their entirety.

If the nucleic acid encoding OBG3 polypeptide fragments lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the insert from the OBG3 polypeptide fragment cDNA lacks a poly A signal, this sequence can be added to the construct by, for example, splicing out the Poly A signal from pSG5 (Stratagene) using BglI and SalI restriction endonuclease enzymes and incorporating it into the mammalian expression vector pXT1 (Stratagene). pXT1 contains the LTRs and a portion of the gag gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex Thymidine Kinase promoter and the selectable neomycin gene.

The nucleic acid encoding an OBG3 fragment can be obtained by PCR from a vector containing the OBG3 nucleotide sequence using oligonucleotide primers complementary to the desired OBG3 cDNA and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and BglIII at the 5' end of the corresponding cDNA 3' primer, taking care to ensure that the sequence encoding the OBG3 fragment is positioned properly with respect to the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1, now containing a poly A signal and digested with BglIII. Alternative methods are presented in Examples 1-3.

Transfection of an OBG3 fragment-expressing vector into mouse NIH 3T3 cells is one embodiment of introducing polynucleotides into host cells. Introduction of a polynucleotide encoding a polypeptide into a host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al. ((1986) *Methods in Molecular Biology*, Elsevier Science Publishing Co., Inc., Amsterdam). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector. Methods of expressing OBG3 fragment of the invention in cells are described in Examples 1-3.

A polypeptide of this invention (i.e. an OBG3 or gOBG3 fragment) can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or

ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells.

Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. Preferably the polypeptides of the invention are non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (*e.g.*, coding sequence), and/or to include genetic material (*e.g.*, heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (*e.g.*, promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination, see, *e.g.*, US Patent Number 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., (1989) *Proc Natl Acad Sci USA* 86(22):8932-5; Koller et al., (1989) *Proc Natl Acad Sci USA* 86(22):8927-31; and Zijlstra et al. (1989) *Nature* 342(6248):435-8; the disclosures of each of which are incorporated by reference in their entireties).

Modifications

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (*See, e.g.*, Creighton, 1983 *Proteins*. New York, New York: W.H. Freeman and Company; and Hunkapiller et al., (1984) *Nature* 310(5973):105-11). For example, a relative short fragment of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution

or addition into the fragment sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, a-amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, g-Abu, c-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, 5 citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoroamino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptide fragments which are differentially modified during 10 or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄; acetylation, formylation, oxidation, reduction; 15 metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends, attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a 20 result of procaryotic host cell expression. The polypeptide fragments may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the polypeptide.

Also provided by the invention are chemically modified derivatives of the polypeptides of the invention that may provide additional advantages such as increased solubility, stability and 25 circulating time of the polypeptide, or decreased immunogenicity. See U.S. Patent No: 4,179,337. The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more 30 attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may 35 be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the polypeptide with consideration of effects on functional or antigenic domains of the polypeptide. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF, see also Malik et al. (1992) Exp Hematol 20(8):1028-35, reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues, glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus may be accomplished by reductive alkylation, which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

In further preferred embodiment, the invention features a method of reducing body mass comprising providing or administering to individuals in need of reducing body mass said pharmaceutical or physiologically acceptable composition described in the fifth aspect in combination with provision or administration of an antagonist of dipeptidyl peptidase cleavage of OBG3 or gOBG3 polypeptide fragment of the first aspect.

Preferred said antagonist is a peptidyl derivative of a diester of alpha-aminoalkylphosphonic acid (US Patent Number 5,543,396 which disclosure is hereby incorporated by reference in its entirety). More preferred said peptidyl derivative is selected from Ala-Pro^P(OZ)₂, AcOH.Ala-Pip^P(Oph)₂, HCl.Ala-Pro^P(Oph-4Cl)₂, HCl.Ala-Pip^P(Oph-4Cl)₂, or 2HCl.Lys-Pip^P(Oph-4Cl)₂, where Z represents an aryl group, a substituted aryl group or a highly fluorinated alkyl group, Pro^P represents a proline phosphonate derivative, and Pip^P represents piperidyl phosphonate (US Patent Number 5,543,396 which disclosure is hereby incorporated by reference in its entirety).

Other preferred said antagonist is a compound of the general formula Z-Xaa-Y', in which Xaa is an amino acid, Z is a protecting group, and Y' is one of various types of ring structures (US Patent Number 6,090,786 which disclosure is hereby incorporated by reference in its entirety). More preferred is said compound wherein Z may or may not be present and represents a protecting group, such as benzyloxycarbonyl; Xaa represents alanine, methionine, arginine, phenylalanine, aspartic acid, proline, asparagine, serine, cysteine, threonine, glycine, tyrosine, glutamic acid, tryptophan, glutamine, valine, isoleucine, lysine, leucine, L-thioproline, L-homoproline, L-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic), L-2,3-dihydroindol-2-carboxylic acid, L-naphthylglycine, L-phenylglycine, L-4-phenylproline, O-benzyl tyrosine, omega-Z lysine, or omega-acetyl lysine; and Y' represents a pyrrolidine, a phosphonate or phosphinate derivative, or reduced peptide; or pharmaceutically acceptable salts thereof (US Patent Number 6,090,786 which disclosure is hereby incorporated by reference in its entirety).

Other preferred said antagonist is sulphostin (US Patent Number 6,214,340 which disclosure is hereby incorporated by reference in its entirety).

Other preferred said antagonist is N-(substituted glycyI)-2-cyanopyrrolidine (US Patent Number 6,166,063). More preferred said N-(substituted glycyI)-2-cyanopyrrolidine is selected from pyrrolidine, 1-[[[(3,5-dimethyl-1-adamantyI)amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-ethyl-1-adamantyI)amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-methoxy-1-adamantyI)amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-[[[(t-butylamino)carbonyl]oxy]-1-adamantyI]amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-[[[(4-methoxyphenyl)amino]-carbonyl]oxy]-1-adamantyI]amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-[[[(phenylamino)carbonyl]oxy]-1-adamantyI]amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(5-hydroxy-2-adamantyI)amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-acetyloxy-1-adamantyI)amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-[[[(diisopropyl)amino]carbonyl]oxy]-1-adamantyI]amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-[[[(cyclohexyl)amino]carbonyl]oxy]-1-adamantyI]amino]-acetyl]-2-cyano-, (S)-; pyrrolidine, 1-[[[(3-ethoxy-1-adamantyI)amino]-acetyl]-2-cyano-, (S)-; or, in each case, a pharmaceutically acceptable acid addition salt thereof (US Patent Number 6,166,063).

Other preferred said antagonist is tetrahydroisoquinoline 3-carboxamide derivative of formula ##STR1## (US Patent Number 6,172,081). More preferred is said derivative and pharmaceutically acceptable salts thereof wherein X is CH₂, S, O, or C(CH₃)₂; R₁ and R₂ are independently hydrogen, hydroxy, alkyl, alkoxy, aralkoxy, or halogen (US Patent Number 6,172,081).

Other preferred said antagonist is valine-pyrrolidide [Deacon (2001) Diabetes 50:1588-1597 which disclosure is hereby incorporated by reference in its entirety].

Multimers

The polypeptide fragments of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers

and multimers of the polypeptide fragments of the invention, their preparation, and compositions (preferably, pharmaceutical or physiologically acceptable compositions) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the OBG3 polypeptide fragments of the invention (including polypeptide fragments, variants, splice variants, and fusion proteins corresponding to these polypeptide fragments as described herein). These homomers may contain polypeptide fragments having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptide fragments having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptide fragments having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing polypeptide fragments having identical or different amino acid sequences) or a homotrimer (e.g., containing polypeptide fragments having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., corresponding to different proteins or polypeptide fragments thereof) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in the sequence listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences, which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent

associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteopontin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins, and have since been found in a variety of different proteins (Landschulz et al., (1988) Genes Dev. Jul;2(7):786-800). Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. FEBS Letters (1994) 344(2-3):191-5 and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention. In another example, proteins of the invention are associated by interactions between Flag® & polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti Flag® antibody.

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C-terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, at least 30 techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (See, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

II. Pharmaceutical or Physiologically Acceptable Compositions of the Invention

The OBG3 and gOBG3 polypeptide fragments of the invention can be administered to non-human animals and/or humans, alone or in pharmaceutical or physiologically acceptable compositions where they are mixed with suitable carriers or excipient(s). The pharmaceutical or physiologically acceptable composition is then provided at a therapeutically effective dose. A therapeutically effective dose refers to that amount of OBG3 or gOBG3 fragment sufficient to result in prevention or amelioration of symptoms or physiological status of obesity-related diseases or disorders as determined

by the methods described herein. A therapeutically effective dose can also refer to the amount of OBG3 or gOBG3 fragment necessary for a reduction in weight or a prevention of an increase in weight or prevention of an increase in the rate of weight gain in persons desiring this affect for cosmetic reasons. A therapeutically effective dosage of an OBG3 or gOBG3 fragment of the invention is that dosage that is adequate to promote weight loss or weight gain with continued periodic use or administration. Techniques for formulation and administration of OBG3 polypeptide fragments may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

Other diseases or disorders that OBG3 polypeptide fragments of the invention could be used to treat or prevent include, but are not limited to, obesity and obesity-related diseases and disorders such as obesity, impaired glucose tolerance, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. Yet other obesity-related diseases or disorders of the invention include cachexia, wasting, AIDS-related weight loss, cancer-related weight loss, anorexia, and bulimia. The OBG3 or gOBG3 polypeptide fragments may also be used to enhance physical performance during work or exercise or enhance a feeling of general well-being. Physical performance activities include walking, running, jumping, lifting and/or climbing.

The OBG3 or gOBG3 polypeptide fragments or antagonists thereof may also be used to treat dyslexia, attention-deficit disorder (ADD), attention-deficit/hyperactivity disorder (ADHD), and psychiatric disorders such as schizophrenia by modulating fatty acid metabolism, more specifically, the production of certain long-chain polyunsaturated fatty acids.

It is expressly considered that the OBG3 or gOBG3 polypeptide fragments of the invention may be provided alone or in combination with other pharmaceutically or physiologically acceptable compounds. Other compounds useful for the treatment of obesity and other diseases and disorders are currently well-known in the art.

In a preferred embodiment, the OBG3 or gOBG3 polypeptide fragments are useful for, and used in, the treatment of insulin resistance and diabetes using methods described herein and known in the art. More particularly, a preferred embodiment relates to process for the therapeutic modification and regulation of glucose metabolism in an animal or human subject, which comprises administering to a subject in need of treatment (alternatively on a timed daily basis) an OBG or OBG3 polypeptide fragment (or polynucleotide encoding said polypeptide) in dosage amount and for a period sufficient to reduce plasma glucose levels in said animal or human subject.

Further preferred embodiments relate to methods for the prophylaxis or treatment of diabetes comprising administering to a subject in need of treatment (alternatively on a timed daily basis) an OBG or OBG3 polypeptide fragment (or polynucleotide encoding said polypeptide) in dosage amount and for a period sufficient to reduce plasma glucose levels in said animal or human subject.

Routes of Administration

The composition according to the present invention is preferably administered systemically. The administration can be carried out by methods familiar to a person skilled in the art, for example, intracisternally, intravenously, or peripherally. For the intracisternal or intravenous administration, OBG3 can be suspended, for example, in physiologic saline. However, infusion or bolus injections at regular time intervals are particularly preferred.

Methods of introduction include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, oral, intrapulmonary, and intranasal administration. In addition, it may be desirable to introduce the pharmaceutical composition of the invention into the central nervous system by any suitable route, including intrathecal, e.g. intraventricular injection. Intraventricular injection may be facilitated by an intraventricular catheter, for example attached to a reservoir, such as an implantable port catheter system.

Furthermore, it may be desirable to administer the pharmaceutical compositions, which are used according to the invention, locally to the area in need of treatment. This may be achieved by, for example, and not by way of limitation, injection by means of a catheter or by means of an implant, said implant being of a porous, non-porous or gelatinous material, including membranes, such as sialastic membranes or fibers.

EXAMPLES

The following Examples are provided for illustrative purposes and not as a means of limitation. One of ordinary skill in the art would be able to design equivalent assays and methods based on the disclosure herein all of which form part of the instant invention.

It should be noted that the term full-length OBG3 polypeptide used throughout the specification is intended to encompass the protein homologs ACRP30 [Scherer, *et al.*, "A novel serum protein similar to C1q, produced exclusively in adipocytes"; *J Biol Chem* 270, 26746-26749 (1995)], AdipoQ [Hu, *et al.*, "AdipoQ is a novel adipose-specific gene dysregulated in obesity", *J Biol Chem* 271, 10697-10703 (1996)] and the human homolog APM1 [Maeda, *et al.*, "cDNA cloning and expression of a novel adipose specific collagen-like factor, APM1 (AdiPose Most abundant Gene transcript 1)", *Biochem Biophys Res Commun* 221, 286-289 (1996)] or GBP28 [Nakano, *et al.*, "Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma", *J Biochem (Tokyo)* 120, 803-812 (1996)]. OBG3 is also intended to encompass other homologs.

EXAMPLE 1: Demyelination in "Gottingen Mini Pigs"

Lesion:

An experimental model of demyelination has been established in the female adult "Gottingen Mini Pig" (age: 10-14 months, weight 25-30 kg) by stereotaxic injection of lysolecithin (LL) at multiple sites into periventricular subcortical white matter of the brain (usually 2-3 injection sites located in one hemisphere; 5 ul of 1% LL in 0.9% saline per lesion infused over a period of 15 min.). This infusion causes a rapid reduction of myeline sheaths within the diffusion area of the detergent (Blakemore, W.F., Neuropathol. Appl. Neurobiol. 4 (1978) 47-59).

OBG3 application:

Immediately after LL-application, a stainless steel needle is implanted in the left ventricle space and affixed with an ionomeric bonding bone cement and an osmotic pump (200 ul volume, infusion rate is connected with the needle and implanted subcutaneously, or a catheter system) is implanted either into the lateral ventricle of the brain or in the subarachnoidal space of the lumer spinal cord.

Ways of application:

1. Chornic infusion of OBG3 via Alzet pump directly intracerebral, beyond the blood brain barrier:

- application interval: 7-20 days
- concentration of OBG3: 0.4mg/ml, dissolved in PBS, pH 7.2-7.4
- dosages: 0.5 ug/h corresponding to 1.2-4 ug/kg/interval

2. Pulsative injections of OBG3:

After fixation of the catheter with bone cement, the intrathecal catheter is connected with a subcutaneous drug delivery system that is implanted subcutaneously in the upper back region. OBG3 (dissolved in phosphate buffer saline) injected every 2nd day through the skin into the port, starting 2 days after initial LL lesions, by single bolus injections into the port at a dose of 0.05-5 ug/kg body weight (3-350 ug/injection). 3-5 injections were carried out into the cerebrospinal fluid within a period of 6-12 days after LL-induced demyelination.

Controls:

Controls are carried out by replacement of OBG3 with an equal dose of Cytochrome C applied in the same ways and time intervals/

Evaluation of OBG3 effects:

At the end of the OBG3 (control, respectively) application period, the experimental animals are sacrificed, the brain removed and prepared for histological examination: the areas of interest were studied with routine histological (H.E. staining, Luxol Fast Blue staining), immunocytochemical (myeline basic protein-MBP) and MRI-analysis.

ADDITIONAL REFERENCES

1. Shapiro, L. & Scherer, P. E., "The crystal structure of a complement-1q family protein suggests an evolutionary link to tumor necrosis factor", *Curr Biol* 8, 335-338 (1998).
- 5 2. Kishore, U. & Reid, K. B., "Modular organization of proteins containing C1q-like globular domain", *Immunopharmacology* 42, 15-21 (1999).
3. Kavety, B. & Morgan, J. I., "Characterization of transcript processing of the gene encoding precerebellin-1", *Brain Res Mol Brain Res* 63, 98-104 (1998).
4. Satoh, F. et al., "Cerebellin and cerebellin mRNA in the human brain, adrenal glands and the
10 tumour tissues of adrenal tumour, ganglioneuroblastoma and neuroblastoma", *J Endocrinol* 154, 27-34 (1997).
5. Mazzocchi, G. et al., "Cerebellin enhances in vitro secretory activity of human adrenal gland", *J Clin Endocrinol Metab* 84, 632-635 (1999).
6. Kondo, N. & Kondo, J., "Identification of novel blood proteins specific for mammalian hibernation", *J Biol Chem* 267, 473-478 (1992).
- 15 7. Takamatsu, N., Ohba, K., Kondo, J., Kondo, N. & Shiba, T., "Hibernation-associated gene regulation of plasma proteins with a collagen-like domain in mammalian hibernators", *Mol Cell Biol* 13, 1516-1521 (1993).
8. Spiegelman, B. M. & Hotamisligil, G. S., "Through thick and thin: wasting, obesity, and
20 TNF alpha", *Cell* 73, 625-627 (1993).
9. Saito, K. et al., "Regulation of gelatin-binding protein 28 (GBP28) gene expression by C/EBP", *Biol Pharm Bull* 22, 1158-1162 (1999).
10. Takamatsu, N. et al., "Expression of multiple alpha1-antitrypsin-like genes in hibernating species of the squirrel family", *Gene* 204, 127-132 (1997).
- 25 11. Ross, S. R., Graves, R. A. & Spiegelman, B. M., "Targeted expression of a toxin gene to adipose tissue: transgenic mice resistant to obesity", *Genes Dev* 7, 1318-1324 (1993).
12. Moitra, J. et al., "Life without white fat: a transgenic mouse", *Genes Dev* 12, 3168-3181 (1998).
13. Peters, S. J., Dyck, D. J., Bonen, A. & Spriet, L. L., "Effects of epinephrine on lipid
30 metabolism in resting skeletal muscle", *Am J Physiol* 275, E300-309 (1998).
14. Rigotti, A., Acton, S. L. & Krieger, M., "The class B scavenger receptors SR-BI and CD36 are receptors for anionic phospholipids", *J Biol Chem* 270, 16221-16224 (1995).
15. Hirsch, D., Stahl, A. & Lodish, H. F., "A family of fatty acid transporters conserved from mycobacterium to man", *Proc Natl Acad Sci U S A* 95, 8625-8629 (1998).
- 35 16. Hotamisligil, G. S. et al., "IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF-alpha- and obesity-induced insulin resistance", *Science* 271, 665-668 (1996).

17. Peraldi, P. & Spiegelman, B., "TNF-alpha and insulin resistance: summary and future prospects", *Mol Cell Biochem* 182, 169-175 (1998).
18. Kirchgessner, T. G., Uysal, K. T., Wiesbrock, S. M., Marino, M. W. & Hotamisligil, G. S., "Tumor necrosis factor-alpha contributes to obesity-related hyperleptinemia by regulating leptin release from adipocytes", *J Clin Invest* 100, 2777-2782 (1997).
19. Uysal, K. T., Wiesbrock, S. M., Marino, M. W. & Hotamisligil, G. S., "Protection from obesity-induced insulin resistance in mice lacking TNF- alpha function", *Nature* 389, 610-614 (1997).
20. Jensen, D. R. et al., "Prevention of diet-induced obesity in transgenic mice overexpressing skeletal muscle lipoprotein lipase", *Am J Physiol* 273, R683-689 (1997).
21. Levak-Frank, S. et al., "Muscle-specific overexpression of lipoprotein lipase causes a severe myopathy characterized by proliferation of mitochondria and peroxisomes in transgenic mice", *J Clin Invest* 96, 976-986 (1995).
22. Scheja, L. et al., "Altered insulin secretion associated with reduced lipolytic efficiency in aP2-/- mice", *Diabetes* 48, 1987-1994 (1999).
23. Griffin, M. E. et al., "Free fatty acid-induced insulin resistance is associated with activation of protein kinase C theta and alterations in the insulin signaling cascade", *Diabetes* 48, 1270-1274 (1999).
24. Dresner, A. et al., "Effects of free fatty acids on glucose transport and IRS-1-associated phosphatidylinositol 3-kinase activity", *J Clin Invest* 103, 253-259 (1999).
25. Kelley, D. E., Goodpaster, B., Wing, R. R. & Simoneau, J. A., "Skeletal muscle fatty acid metabolism in association with insulin resistance, obesity, and weight loss", *Am J Physiol* 277, E1130-1141 (1999).
26. Roden, M. et al., "Mechanism of free fatty acid-induced insulin resistance in humans", *J Clin Invest* 97, 2859-2865 (1996).
27. Deacon, C. F. et al., "Dipeptidyl peptidase IV inhibition reduces the degradation and clearance of GLP and potentiates its insulinotropic and antihyperglycemic effects in anesthetized pigs", *Diabetes* 50, 1588-1597 (2001).

CLAIMS

What is claimed is:

- 5 1. A method of increasing the rate of regeneration of myelin sheaths of axons comprising the step of contacting an oligodendrocyte progenitor cell with a composition comprising a polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 6 or a biologically active fragment thereof.
- 10 2. A method of increasing the rate of cellular division of oligodendrocyte progenitor cells in which said oligodendrocyte progenitor cells are contacted with a composition comprising a polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 6 or a biologically active fragment thereof.
- 15 3. A method of increasing the rate of migration of oligodendrocyte progenitor cells from the germinal centers to demyelinated axons in the CNS in which said oligodendrocyte progenitor cells are contacted with a composition comprising a polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 6 or a biologically active fragment thereof.
- 20 4. A method of inducing oligodendrocyte progenitor cells to differentiate into mature dendrocytic cells at sites of demyelination in axons in the CNS in which said oligodendrocyte progenitor cells are contacted with a composition comprising a polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 6 or a biologically active fragment thereof.
- 25 5. The method of any one of claims 1-4, wherein said method is used to treat multiple sclerosis.
6. The method of any one of claims 1-4, wherein said method is used to treat hereditary leukodystrophies selected from the group consisting of metachromatic leukodystrophy, Refsum's disease, adrenoleukodystrophy, Krabbe's disease, phenylketonuria, Canavan disease, Pelizaeus-Merzbacher disease, and Alexander's disease.
- 30 7. The method of any one of claims 1-4, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of amino acids 84-244, 85-244, 101 to 244, 102-244, or 103-244 of SEQ ID NO:6.
8. The method of any one of claims 1-4, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of amino acids 166-193, amino acids 166-176, or amino acids 167-176 of SEQ ID NO: 6 or a biologically active fragment thereof.
- 35 9. The use of a pharmaceutical composition for the treatment of diseases in which demyelination of nerve fibers occurs, comprising an amino acid sequence selected from the group consisting of amino acids 84-244, 85-244, 101 to 244, 102-244, 103-244, 166-193, 166-176, or 167-176 of SEQ ID NO:6 or a biologically active fragment thereof.
10. A method for the production of a pharmaceutical composition for use in the treatment of disease in which demyelination of nerve fibers occurs, characterized in that an amino

acid sequence selected from the group consisting of amino acids 84-244, 85-244, 101 to 244, 102-244, 103-244, 166-193, 166-176, or 167-176 of SEQ ID NO:6 or a biologically active fragment thereof is used as an essential component of the composition.

11. The method of any one of claims 1-4, 7 or 8, wherein said polypeptide is
5 administered to an individual in need thereof.

SEQUENCE LISTING

<110> GENSET

<120> Use of OBG3 for Promoting Central Nervous System Remyelination

<130> 161.W01

<150> 60/332,119

<151> 2001-11-21

<160> 7

<170> Patent.pm

<210> 1

<211> 1152

<212> DNA

<213> mus musculus

<400> 1

```

gaattcgcca cgaggg atg cta ctg ttg caa gct ctc ctg ttc ctc tta atc      52
              Met Leu Leu Leu Gln Ala Leu Leu Phe Leu Leu Ile
              1              5              10
ctg ccc agt cat gcc gaa gat gac gtt act aca act gaa gag cta gct      100
Leu Pro Ser His Ala Glu Asp Asp Val Thr Thr Thr Glu Glu Leu Ala
              15              20              25
cct gct ttg gtc cct cca ccc aag gga act tgt gca ggt tgg atg gca      148
Pro Ala Leu Val Pro Pro Pro Lys Gly Thr Cys Ala Gly Trp Met Ala
              30              35              40
ggc atc cca gga cat tct ggc cac aat ggc aca cca ggc cgt gat ggc      196
Gly Ile Pro Gly His Ser Gly His Asn Gly Thr Pro Gly Arg Asp Gly
              45              50              55              60
aga gat ggc act cct gga gag aag gga gag aaa gga gat tca ggt ctt      244
Arg Asp Gly Thr Pro Pro Gly Glu Lys Gly Glu Lys Gly Asp Ser Gly Leu
              65              70              75
ctt ggt cct aag ggt gag aca gga gat gtt gga atg aca gga gct gaa      292
Leu Gly Pro Lys Gly Glu Thr Gly Asp Val Gly Met Thr Gly Ala Glu
              80              85              90
ggg cct cgg ggc ttc ccc gga acc cct ggc agg aaa gga gag cct gga      340
Gly Pro Arg Gly Phe Pro Gly Thr Pro Gly Arg Lys Gly Glu Pro Gly
              95              100              105
gaa gcc gct tat gtg tat cgc tca ggc ttc agt gtg ggg ctg gag acc      388
Glu Ala Ala Tyr Val Tyr Arg Ser Gly Phe Ser Val Gly Leu Glu Thr
              110              115              120
cgc gtc act gtt ccc aat gta ccc att cgc ttt act aag atc ttc tac      436
Arg Val Thr Val Pro Asn Val Pro Ile Arg Phe Thr Lys Ile Phe Tyr
              125              130              135              140
aac caa cag aat cat tat gac aac agc act ggc aag ttc tac tgc aac      484
Asn Gln Gln Asn His Tyr Asp Asn Ser Thr Gly Lys Phe Tyr Cys Asn
              145              150              155
att ccg gga ctc tac ttc tct tac cac atc acg gtg tac atg aaa      532
Ile Pro Gly Leu Tyr Tyr Phe Ser Tyr His Ile Thr Val Tyr Met Lys
              160              165              170
gat gtg aag gtg agc ctc ttc aag aag gac aag gcc gtt ctc ttc acc      580
Asp Val Lys Val Ser Leu Phe Lys Lys Asp Lys Ala Val Leu Phe Thr
              175              180              185
tac gac cag tat cag gaa aag aat gtg gac cag gcc tct ggc tct gtg      628
Tyr Asp Gln Tyr Gln Glu Lys Asn Val Asp Gln Ala Ser Gly Ser Val
              190              195              200
ctc ctc cat ctg gag gtg gga gac caa gtc tgg ctc cag gtg tat ggg      676
Leu Leu His Leu Glu Val Gly Asp Gln Val Trp Leu Gln Val Tyr Gly
              205              210              215              220

```

gat ggg gac cac aat gga ctc tat gca gat aac gtc aac gac tct aca	724
Asp Gly Asp His Asn Gly Leu Tyr Ala Asp Asn Val Asn Asp Ser Thr	
225 230 235	
ttt act ggc ttt ctt ctc ttc cat gat acc aac tga ctgcaactac	770
Phe Thr Gly Phe Leu Leu Phe His Asp Thr Asn *	
240 245	
tcataagccca tacaccagga gaatcatgga acgtcgacac actttcagct tagtttgaga	830
gattgatttt attgcttagt ttgagagtc tgagtattat ccacacgtgt actcacttgt	890
tcattaaacg actttataaa aaataatttg tgttcttagt ccagaaaaaa aggcactccc	950
tggtctccac gactcttaca tggtagcaat aacagaatga aaatcacatt tggtagggg	1010
gcttcacaat attcgcatga ctgtctggaa gtagaccatg ctattttct gctcactgta	1070
cacaaatatt gtccacataa accctataat gtaaatatga aatacagtga ttattctctc	1130
aaaaaaaaa cgtgcgcaat tc	1152

<210> 2

<211> 247

<212> PRT

<213> mus musculus

<400> 2

Met Leu Leu Leu Gln Ala Leu Leu Phe Leu Leu Ile Leu Pro Ser His	
1 5 10 15	
Ala Glu Asp Asp Val Thr Thr Thr Glu Glu Leu Ala Pro Ala Leu Val	
20 25 30	
Pro Pro Pro Lys Gly Thr Cys Ala Gly Trp Met Ala Gly Ile Pro Gly	
35 40 45	
His Ser Gly His Asn Gly Thr Pro Gly Arg Asp Gly Arg Asp Gly Thr	
50 55 60	
Pro Gly Glu Lys Gly Glu Lys Gly Asp Ser Gly Leu Leu Gly Pro Lys	
65 70 75 80	
Gly Glu Thr Gly Asp Val Gly Met Thr Gly Ala Glu Gly Pro Arg Gly	
85 90 95	
Phe Pro Gly Thr Pro Gly Arg Lys Gly Glu Pro Gly Glu Ala Ala Tyr	
100 105 110	
Val Tyr Arg Ser Gly Phe Ser Val Gly Leu Glu Thr Arg Val Thr Val	
115 120 125	
Pro Asn Val Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn	
130 135 140	
His Tyr Asp Asn Ser Thr Gly Lys Phe Tyr Cys Asn Ile Pro Gly Leu	
145 150 155 160	
Tyr Tyr Phe Ser Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val	
165 170 175	
Ser Leu Phe Lys Lys Asp Lys Ala Val Leu Phe Thr Tyr Asp Gln Tyr	
180 185 190	
Gln Glu Lys Asn Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu	
195 200 205	
Glu Val Gly Asp Gln Val Trp Leu Gln Val Tyr Gly Asp Gly Asp His	
210 215 220	
Asn Gly Leu Tyr Ala Asp Asn Val Asn Asp Ser Thr Phe Thr Gly Phe	
225 230 235 240	
Leu Leu Phe His Asp Thr Asn	
245	

<210> 3

<211> 1276

<212> DNA

<213> mus musculus

<400> 3

ctctaaagat tgtcagtgga tctgacgaca ccaaaagggc tcagg atg cta ctg ttg	57
Met Leu Leu Leu	
1	
caa gct ctc ctg ttc ctc tta atc ctg ccc agt cat gcc gaa gat gac	105

Gln Ala Leu Leu Phe Leu Ile Leu Pro Ser His Ala Glu Asp Asp
 5 10 15 20
 gtt act aca act gaa gag cta gct cct gct ttg gtc cct cca ccc aag 153
 Val Thr Thr Thr Glu Leu Ala Pro Ala Leu Val Pro Pro Lys
 25 30 35
 gga act tgt gca ggt tgg atg gca ggc atc cca gga cat cct ggc cac 201
 Gly Thr Cys Ala Gly Trp Met Ala Gly Ile Pro Gly His Pro Gly His
 40 45 50
 aat ggc aca cca ggc cgt gat ggc aga gat ggc act cct gga gag aag 249
 Asn Gly Thr Pro Gly Arg Asp Gly Arg Asp Gly Thr Pro Gly Glu Lys
 55 60 65
 gga gag aaa gga gat gca ggt ctt ctt ggt cct aag ggt gag aca gga 297
 Gly Glu Lys Gly Asp Ala Gly Leu Leu Gly Pro Lys Gly Glu Thr Gly
 70 75 80
 gat gtt gga atg aca gga gct gaa ggg cca cgg ggc ttc ccc gga acc 345
 Asp Val Gly Met Thr Gly Ala Glu Gly Pro Arg Gly Phe Pro Gly Thr
 85 90 95 100
 cct ggc agg aaa gga gag cct gga gaa gcc gct tat atg tat cgc tca 393
 Pro Gly Arg Lys Gly Glu Pro Gly Glu Ala Tyr Met Tyr Arg Ser
 105 110 115
 gcg ttc agt gtg ggg ctg gag acc cgc gtc act gtt ccc aat gta ccc 441
 Ala Phe Ser Val Gly Leu Glu Thr Arg Val Thr Val Pro Asn Val Pro
 120 125 130
 att cgc ttt act aag atc ttc tac aac caa cag aat cat tat gac ggc 489
 Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn His Tyr Asp Gly
 135 140 145
 agc act ggc aag ttc tac tgc aac att cgc gga ctc tac tac ttc tct 537
 Ser Thr Gly Lys Phe Tyr Cys Asn Ile Pro Gly Leu Tyr Tyr Phe Ser
 150 155 160
 tac cac atc acg gtg tac atg aaa gat gtg aag gtg agc ctc ttc aag 585
 Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val Ser Leu Phe Lys
 165 170 175 180
 aag gac aag gcc gtt ctc ttc acc tac gac cag tat cag gaa aag aat 633
 Lys Asp Lys Ala Val Leu Phe Thr Tyr Asp Lys Gln Glu Lys Asn
 185 190 195
 gtg gac cag gcc tct ggc tct gtg ctc ctc cat ctg gag gtg gga gac 681
 Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu Glu Val Gly Asp
 200 205 210
 caa gtc tgg ctc cag gtg tat ggg gat ggg gac cac aat gga ctc tat 729
 Gln Val Trp Leu Gln Val Tyr Gly Asp Gly Asp His Asn Gly Leu Tyr
 215 220 225
 gca gat aac gtc aac gac tct aca ttt act ggc ttt ctt ctc tac cat 777
 Ala Asp Asn Val Asn Asp Ser Thr Phe Thr Gly Phe Leu Leu Tyr His
 230 235 240
 gat acc aac tga ctgcaactac ccatagccca tacaccagga gaatcatgga 829
 Asp Thr Asn *
 245
 acagtcgaca cactttcagc ttagtttgag agattgattt tattgcttag tttgagagtc 889
 ctgagtattca tccacacgtg tactcacttg ttcattaaac gactttataa aaaataattt 949
 gtgttctctag tccagaaaaa aaggcactcc ctggtctcca cgactottac atggtagcaa 1009
 taacagaatg aaaaatcacat ttgggtatggg ggcttcacaa tattcgcgat actgtctgga 1069
 agtagacat gctatttttc tgcctactgt acacaaatat tgttcacata aaccctataa 1129
 tgtaaatatg aaatacagtg attactcttc tcacaggctg agtgtatgaa tgtctaaaga 1189
 cccataagta ttaaagtggg agggataaat tggaaaaaaa aaaaaaaa aagaaaaact 1249
 ctgagcaca ctggcggcgg ttactag 1276

<210> 4

<211> 247

<212> PRT

<213> mus musculus

<400> 4

Met Leu Leu Leu Gln Ala Leu Leu Phe Leu Leu Ile Leu Pro Ser His

4

1	5	10	15
Ala Glu Asp Asp Val Thr Thr Thr Glu Glu Leu Ala Pro Ala Leu Val			
Pro Pro Pro Lys Gly Thr Cys Ala Gly Trp Met Ala Gly Ile Pro Gly	20	25	30
His Pro Gly His Asn Gly Thr Pro Gly Arg Asp Gly Arg Asp Gly Thr	35	40	45
Pro Gly Glu Lys Gly Glu Lys Gly Asp Ala Gly Leu Leu Gly Pro Lys	50	55	60
Gly Glu Thr Gly Asp Val Gly Met Thr Gly Ala Glu Gly Pro Arg Gly	65	70	75
Phe Pro Gly Thr Pro Gly Arg Lys Gly Glu Pro Gly Glu Ala Ala Tyr	85	90	95
Met Tyr Arg Ser Ala Phe Ser Val Gly Leu Glu Thr Arg Val Thr Val	100	105	110
Pro Asn Val Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn	115	120	125
His Tyr Asp Gly Ser Thr Gly Lys Phe Tyr Cys Asn Ile Pro Gly Leu	130	135	140
Tyr Tyr Phe Ser Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val	145	150	155
Ser Leu Phe Lys Lys Asp Lys Ala Val Leu Phe Thr Tyr Asp Gln Tyr	165	170	175
Gln Glu Lys Asn Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu	180	185	190
Glu Val Gly Asp Gln Val Trp Leu Gln Val Tyr Gly Asp Gly Asp His	195	200	205
Asn Gly Leu Tyr Ala Asp Asn Val Asn Asp Ser Thr Phe Thr Gly Phe	210	215	220
Leu Leu Tyr His Asp Thr Asn	225	230	235
	245		240

<210> 5

<211> 4517

<212> DNA

<213> Homo sapiens

<400> 5

ctgattccat accagagggg ctcagg atg ctg ttg ctg gga gct gtt cta ctg	53
Met Leu Leu Leu Gly Ala Val Leu Leu	
cta tta gct ctg ccc ggg cat gac cag gaa acc acg act caa ggg ccc	101
Leu Leu Ala Leu Pro Gly His Asp Gln Glu Thr Thr Gln Gly Pro	
gga gtc ctg ctt ccc ctg ccc aag ggg gcc tgc aca ggt tgg atg gcg	149
Gly Val Leu Leu Pro Leu Pro Lys Gly Ala Cys Thr Gly Trp Met Ala	
ggc atc cca ggg cat cgc ggc cat aat ggg gcc cca ggc cgt gat ggc	197
Gly Ile Pro Gly His Pro Gly His Asn Gly Ala Pro Gly Arg Asp Gly	
aga gat ggc acc cct ggt gag aag ggt gag aaa gga gat cca ggt ctt	245
Arg Asp Gly Thr Pro Gly Glu Lys Gly Glu Lys Gly Asp Pro Gly Leu	
att ggt cct aag gga gac atc ggt gaa acc gga gta ccc ggg gct gaa	293
Ile Gly Pro Lys Gly Asp Ile Gly Glu Thr Gly Val Pro Gly Ala Glu	
ggt ccc cga ggc ttt cgc gga atc caa ggc agg aaa gga gaa cct gga	341
Gly Pro Arg Gly Phe Pro Gly Ile Gln Gly Arg Lys Gly Glu Pro Gly	
gaa ggt gcc tat gta tac cgc tca gca ttc agt gtg gga ttg gag act	389
Glu Gly Ala Tyr Val Tyr Arg Ser Ala Phe Ser Val Gly Leu Glu Thr	
tac gtt act atc ccc aac atg ccc att cgc ttt acc aag atc ttc tac	437

[illegible]

```

tgttttctg  tgatgccaa  agaggagaga  ggaagctctc  tttccagat  gcccagcaa  3231
gtgtaacctt  gcattctcatt  gctctggctg  agttgtgtgc  ctgtttctga  ccaatcactg  3291
agtcaaggagg  atgaaatatt  catattgact  taattgcagc  ttaagttagg  ggtatgtaga  3351
gggtattttc  ctaaagcaaa  atggggacac  tgtttacaga  aataggagag  tggatgatag  3411
atgcacaaata  atacctgtcc  acaacaaact  cttaatgctg  tgtttgagct  ttcattgagtt  3471
tcccagagag  acatagctgg  aaaatttcta  ttgattttct  ctaaaatttc  aacaagtatg  3531
taaagtctgg  ctatgctcac  agtctcacat  ctgggtgggg  tgggctcctt  acagaacacg  3591
ctttcacagt  taccctaaac  tctctggggc  agggttattc  ctttgggaa  ccagaggcac  3651
agagacagtc  aactgaggcc  caacagaggg  ctgagagaaa  ctgaggtcaa  gatttcagga  3711
ttaatggctc  tgtgatgctt  tgaagtacaa  ttgtggattt  gtccaattct  ctttagttct  3771
gtcagctttt  gcttcataata  ttttagcgct  ctattattag  atatacatat  gtttagtatt  3831
atgtcttatt  ggtgcattta  ctctcttatt  attatgtaat  gtcccttctt  atctgtgata  3891
attttctgtg  tttcgaagtc  tactttgtct  aaaaataaca  taogcactca  accttctctt  3951
ctttctctct  tctcttcttt  ctctctctct  tttcttctct  ctctctcttt  ccttctcttc  4011
tctctctctt  tctctctctc  tctctctctc  tctctctctc  tgacagactc  tctgtctgtg  4071
gcctcgctg  gagttcagtg  gtgtgatctt  ggctcactgc  tactctacc  atgagcaatt  4131
ctctcgctc  agctcccaa  gtatgctgaa  ctacaggctc  atgccaactg  gccagctaa  4191
tttttgattt  ttttgtagtg  acggggtttc  accacattcg  tcaggttggt  ttcaaaactc  4251
tgactttgtg  atccaccgc  ctccgctctc  caaagtgtcg  ggattacagg  catgagcat  4311
cacaccgtgc  caactttctt  ttgattagtg  tttttgtggt  atatcttttt  coactcatggt  4371
actttaaata  tatctatatt  attgtattta  aaatgtgttt  cttacagact  gcattgagtt  4431
gggtataatt  tttatccagt  ctaaaaaaat  ctgtctttta  attggtgttt  agacaattta  4491
tatttaataa  aatggtggaa  ttttaa

```

<210> 6
 <211> 244
 <212> PRT
 <213> Homo sapiens

```

<400> 6
Met Leu Leu Leu Gly Ala Val Leu Leu Leu Leu Ala Leu Pro Gly His
1      5      10      15
Asp Gln Glu Thr Thr Thr Gln Gly Pro Gly Val Leu Leu Pro Leu Pro
20     25     30
Lys Gly Ala Cys Thr Gly Trp Met Ala Gly Ile Pro Gly His Pro Gly
35     40     45
His Asn Gly Ala Pro Gly Arg Asp Gly Arg Asp Gly Thr Pro Gly Glu
50     55     60
Lys Gly Glu Lys Gly Asp Pro Gly Leu Ile Gly Pro Lys Gly Asp Ile
65     70     75     80
Gly Glu Thr Gly Val Pro Gly Ala Glu Gly Pro Arg Gly Phe Pro Gly
85     90     95
Ile Gln Gly Arg Lys Gly Glu Pro Gly Glu Gly Ala Tyr Val Tyr Arg
100    105    110
Ser Ala Phe Ser Val Gly Leu Glu Thr Tyr Val Thr Ile Pro Asn Met
115    120    125
Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn His Tyr Asp
130    135    140
Gly Ser Thr Gly Lys Phe His Cys Asn Ile Pro Gly Leu Tyr Tyr Phe
145    150    155    160
Ala Tyr His Ile Thr Val Tyr Met Lys Asp Val Lys Val Ser Leu Phe
165    170    175
Lys Lys Asp Lys Ala Met Leu Phe Thr Tyr Asp Gln Tyr Gln Glu Asn
180    185    190
Asn Val Asp Gln Ala Ser Gly Ser Val Leu Leu His Leu Glu Val Gly
195    200    205
Asp Gln Val Trp Leu Gln Val Tyr Gly Glu Gly Glu Arg Asn Gly Leu
210    215    220
Tyr Ala Asp Asn Asp Asn Asp Ser Thr Phe Thr Gly Phe Leu Leu Tyr
225    230    235    240
His Asp Thr Asn

```

<210> 7

<211> 20966
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 1..4811
<223> 5' regulatory region

<220>
<221> exon
<222> 4812..4851
<223> exon 1

<220>
<221> exon
<222> 15144..15365
<223> exon 2

<220>
<221> exon
<222> 16277..20559
<223> exon 3

<220>
<221> misc_feature
<222> 20560..20966
<223> 3' regulatory region

<220>
<221> allele
<222> 3787
<223> 9-27-261 : polymorphic base G or C

<220>
<221> allele
<222> 11118
<223> 99-14387-129 : polymorphic base A or C

<220>
<221> allele
<222> 15120
<223> 9-12-48 : polymorphic base C or T

<220>
<221> allele
<222> 15196
<223> 9-12-124 : polymorphic base G or T

<220>
<221> allele
<222> 15427
<223> 9-12-355 : polymorphic base G or T

<220>
<221> allele
<222> 15500
<223> 9-12-428 : polymorphic base A or G

<220>
<221> allele
<222> 15863
<223> 99-14405-105 : polymorphic base A or G

<220>
<221> allele
<222> 17170
<223> 9-16-189 : polymorphic base deletion of A

<220>
<221> primer_bind
<222> 3528..3545
<223> 9-27.pu

<220>
<221> primer_bind
<222> 3928..3946
<223> 9-27.rp complement

<220>
<221> primer_bind
<222> 10990..11008
<223> 99-14387.pu

<220>
<221> primer_bind
<222> 11423..11442
<223> 99-14387.rp complement

<220>
<221> primer_bind
<222> 15073..15092
<223> 9-12.pu

<220>
<221> primer_bind
<222> 15503..15520
<223> 9-12.rp complement

<220>
<221> primer_bind
<222> 15759..15776
<223> 99-14405.pu

<220>
<221> primer_bind
<222> 16191..16211
<223> 99-14405.rp complement

<220>
<221> primer_bind
<222> 16982..17001
<223> 9-16.pu

<220>
<221> primer_bind
<222> 17384..17402
<223> 9-16.rp complement

<220>
<221> misc_binding
<222> 3775..3799
<223> 9-27-261.probe

<220>
<221> misc_binding


```
<222> 11106..11130
<223> 99-14387-129.probe

<220>
<221> misc_binding
<222> 15108..15132
<223> 9-12-48.probe

<220>
<221> misc_binding
<222> 15184..15208
<223> 9-12-124.probe

<220>
<221> misc_binding
<222> 15415..15439
<223> 9-12-355.probe

<220>
<221> misc_binding
<222> 15488..15512
<223> 9-12-428.probe

<220>
<221> misc_binding
<222> 15851..15875
<223> 99-14405-105.probe

<220>
<221> misc_binding
<222> 17158..17182
<223> 9-16-189.probe

<220>
<221> primer_bind
<222> 3768..3786
<223> 9-27-261.mis

<220>
<221> primer_bind
<222> 3788..3806
<223> 9-27-261.mis complement

<220>
<221> primer_bind
<222> 11099..11117
<223> 99-14387-129.mis

<220>
<221> primer_bind
<222> 11119..11137
<223> 99-14387-129.mis complement

<220>
<221> primer_bind
<222> 15101..15119
<223> 9-12-48.mis

<220>
<221> primer_bind
<222> 15121..15139
<223> 9-12-48.mis complement
```

```

<220>
<221> primer_bind
<222> 15177..15195
<223> 9-12-124.mis

<220>
<221> primer_bind
<222> 15197..15215
<223> 9-12-124.mis complement

<220>
<221> primer_bind
<222> 15408..15426
<223> 9-12-355.mis

<220>
<221> primer_bind
<222> 15428..15446
<223> 9-12-355.mis complement

<220>
<221> primer_bind
<222> 15481..15499
<223> 9-12-428.mis

<220>
<221> primer_bind
<222> 15501..15519
<223> 9-12-428.mis complement

<220>
<221> primer_bind
<222> 15844..15862
<223> 99-14405-105.mis

<220>
<221> primer_bind
<222> 15864..15882
<223> 99-14405-105.mis complement

<220>
<221> primer_bind
<222> 17151..17169
<223> 9-16-189.mis

<220>
<221> primer_bind
<222> 17171..17189
<223> 9-16-189.mis complement

<400> 7
gctgatctgc tgcctcagcc ttcccaaaagt gctgtaattt attaggcata agccactgtg      60
cctgcctagt gttgtacatt ctgtgggttt tgacaattgt atgcatctac atgtatgtac      120
catttatagt attcctgttt ttaatttttag ccattctagt aggcattgtg tgatatctca      180
tggtgattttt aatttgcggtt tcogtaatgg ttaataatgc tgaacatctt tgcattgtgct      240
tgtttgcatt ttgtgtttcc tacttggtga aataattggt catgtcccttt gtccattttc      300
taattgaatt tttttttacc atttagtttt gagatttctt tatacaatct agatccaaat      360
ctcttgctct aaatatgggtt tgcaaataca ttctcttaat tcataatatt ccttttctct      420
ctcttaacag gatgtttcac agagcaaaag ttttagtttt gttgaaatct cacttttcat      480
ttttttcttt agtggattgt gcttttgttg tcatatgtaa gaactcttca ctggccctag      540
atccttgatg tggtttccta agattgccat agcaaatcac catgaactta gtgacaaaaa      600
gacagaaatt tattttcact tctactgtg ggcagactag acgttaatta ttttcatgta      660
tgctcattcc tatgacatct ttctgatata ataattatag ttattcttaa gcttcaccc      720

```

ttttttctatt	agcttttgta	ccttgggtgt	cactttttct	tttttgacat	tgtgacctat	780
gcagatcatc	gtctgttagt	acttagccct	ctccataact	ctccataact	ctctttgtat	840
tcctggagct	tgatgcctga	aatgacacat	cctacattcc	ttggccagat	gggtaccagt	900
tagcttgtgc	acatggggga	caaccgtgaa	aagactgaag	ttgggaagaa	gggagagagt	960
gttgtgttct	agtgagcgcc	cttggcagtg	goggtgacag	tggtcctctg	tcagtgccaa	1020
tggtggagca	gctagcaaga	catgcagtaa	gcgcaggtgc	ataggctatg	gtccaggagc	1080
agtcaccgat	tccttgcttc	taggcaatat	catctccctt	tgctttccca	gcctttctaa	1140
aattatttga	ccttgactag	tacaattttt	tagtatgggg	ggtagtccaa	ggacacaggg	1200
tttaaaaagt	atgaattcag	gggtgcctac	ctgcattgac	tgcgcttgaa	tcagtatggc	1260
ctcttggtcg	gtggcaggag	gtgacagtcc	aaatcatgca	gtagcaaac	agataactaa	1320
attatcatct	gagatacttc	agaagtacag	cogtagccat	actttcagaa	gagatagaga	1380
aattgtctcc	tgggcaggcg	cgggtggctca	cgctgtcat	tcacgacatt	tgaggagccg	1440
agggggtgga	tcacctgagg	tcgggagttc	gagacagcc	tgaccaaac	gggggaacc	1500
tgtctctact	aaaaatacaa	aattagcggg	gcgtggtggc	acatgcccat	aatccagct	1560
actcggggag	ctagggcagg	ataatcgctt	gaacctgaga	ggcagagggt	gcgggtgaact	1620
gagatcatgc	catagtactc	cagcctgggc	aacaagagtg	aaactccatc	tcaaaaaaaa	1680
aaaaaagaaa	aaagaataaa	gaaatgttct	cctttctctg	catttctag	gggtttggga	1740
tggtctacat	gtctgcaggg	cgtgctcact	ctaccaactc	gtcccaactc	tattttttca	1800
aaatcacagt	cttatgcttg	gttacttcag	tttaagattat	ttttaaaaat	cataattaa	1860
caaaaatata	tggtcactgc	taaacatatt	taagtatttt	taagtatttt	ggcctgttct	1920
agtatcccaa	ctcacatgct	aacagggggt	tgacctgtag	ctacggtacc	ctggaggaaa	1980
ctagtcgcat	tattttggta	ttctggctta	agtagtaata	gttctgtctc	gggaaaaaga	2040
tgagctccaa	ggactttctg	attgaatggt	tttcaattac	agctttttaa	ccagtatggc	2100
acagaaactg	ctcttccac	atgacggcct	ttgtgggtgg	tggcagcatt	ccctgagggc	2160
tcgcaaaatg	ctaggctttc	acaatgtcac	tgactgacag	ccaggccagc	caacagtctg	2220
gtgtgattgt	ggggctaaag	tattccaccc	ttgtgcaata	gtcacaagct	ctcttaacca	2280
gctgcattct	tataaagtta	gaagaaaata	cttttttttt	tttgagatgt	attctcgctc	2340
tggtgcccag	gctggagttg	aatgggtcga	tctcggtcgc	ctgcaacctc	cgctcctcgg	2400
gtccaaagca	ttctcctccc	tcagaccccc	gagt-agctgg	gattgcaggt	gcttgccacc	2460
agccccgggt	aaactttttg	tatttttagt	ggagacgggg	tttccacatc	ttcgtcaggg	2520
tggtctcaga	ctctcgactc	caagtgtatc	gcccgcctca	gctccccaaa	atgtcgggat	2580
tacaggcatc	agctactggt	cccggccaaa	gaaaaatact	tttatgcaga	ccctgaaact	2640
acctgaagcg	acatacatca	accttgaggc	ctcacactcc	atcaagaggg	gtgaaaggca	2700
tgaggaaatc	gaagactcag	ggatttttag	ttagacagat	gggtttcaaa	ctctagactt	2760
gtgccttgaa	caaatatttt	acctcattg	aaacttagat	tcattatttg	taaaatgaaa	2820
gacaaataat	gttatctcca	aaggaaagtt	gaatatgata	attcatttat	tcattaatc	2880
aacatttatt	attgctactc	ttgtgccagg	ttctattcta	ggaactaagg	gatacaactt	2940
tgaataggca	aaactctctg	ttctcctga	tttacttttt	tttttttttt	ttgagacaga	3000
gttctcaact	gtgcacccag	gctggagcgc	aatgggtgct	ttggctcact	gcaacctcca	3060
cctcctgggt	tcaagtgatt	ctctgtctct	agcctcccaa	gtagctggga	ctacaggat	3120
gtgccaccac	gcgcggctat	ttctgcattt	ttagttagaga	tggggtttca	catagtgtgc	3180
cagacttgct	tcacaaactc	gatctcaggt	gatattcgct	tcttggtcct	ccaaagtaact	3240
gggattacag	cctgagccca	ctgcacctga	cctgaagttt	atgttctatt	aatgaccaa	3300
agacagtaac	ataaacccaa	aataaatagg	aaaacacat	aaacaaaatc	aaacagtgat	3360
ataattgaga	gttgccttca	tttctttttg	ttgtctcttt	gggttcaatc	gctctgctaa	3420
ctatatggaa	ctctattttc	atggggcaact	tatttaaagc	gggggacact	ggaaagctct	3480
tcagtctctc	catctcaagc	gctcaatgtg	acttctcttg	aaatatattg	acattagcag	3540
gaagctgagg	ctttacatca	gatctttact	ttaatgggtg	acttgacttt	actggttagat	3600
ttttgggtct	gtgtggaact	gtggagatga	tattctgggg	gcaggcagac	acttgccctg	3660
cctctgtctg	agaaaattct	gttttgtagt	ttctgttgaa	gttggtgctg	gctctctaa	3720
ccctgtgctg	ggctgtaatt	taattcatca	gaatgtgtg	cttgcaagaa	ccgctcaga	3780
ttctgcctct	caaaaacaaa	acatgagcgt	gccaaagaa	tccaaggtgt	tgaattgtgc	3840
cacttccagc	ctaaactttc	taggaaaccc	taagtgggtg	gcagcttcca	gttctccagg	3900
ctctctctag	gcacagagct	gggtccacaa	gagacagaa	agggcatat	atgcttaagg	3960
aactggaaaa	acaggctctc	ttctctccac	aaacacacac	acacacatc	acaggttagc	4020
gtcaaaatgt	tatccgaatt	tttggaaaca	aaaaatcttg	aaagatggta	ttccaatctc	4080
acatttttag	taagttttct	attatatagg	attcaaatca	cgattcgagg	cccaactgct	4140
taagaattca	ggggcctttt	aacttgccaa	gccccacacc	actccaggaa	cttccccaca	4200
ccccagttct	cagaattctc	gtgcaaggct	tttctctaat	ccagggtcca	gtccagagag	4260
tgaggatgtg	gctcttaact	ttacctgatt	gcagacccct	ctgacagtgc	ttcctcttga	4320
agcaactcact	gtctgaagct	acacagctct	agacttaact	atgcacagtg	agcaagagct	4380
tggtgtgata	attggcgctc	ctgacttatt	agggcaaatc	tatgggaggg	ggagacactc	4440
tggaccactg	agcaattaat	tcattttacat	taggaagttt	ttccctgcaga	tgcaggaaaa	4500

aaatcttgtt	ttcctgctgt	ggttttgact	tttgccocat	cttctgttgc	tgttgttaga	4560
ggcaaaataa	gggtcaagcc	ctggaacac	aagtgtcttg	actgaagctc	cacttggctt	4620
cgcaagccca	agctgggttg	taccaggttc	cctagggtgc	aggctgtggg	caactgccag	4680
ggacatgttg	ctgccaccgc	gcctctggcc	ctcactgagt	tgcccaattg	gaaatgcaca	4740
ttgtgaggtc	gggaactgcct	gcccccgtga	gtaccaggtc	gttgaggctg	ggccactctc	4800
ttcctacttc	cattctgact	gcagctctgt	gttctgattc	cataccagag	ggtaagagca	4860
attctgtgaa	gttccaggtc	gggtggggga	tgcattgcata	gcctctggct	gggtctaccc	4920
agggtctccc	gtccgttagta	gtgtgggagt	ggatcacagat	ggatactctg	gtccagagcag	4980
cactgggtga	ggcagatatt	cactgggctt	cttccctcgt	tctcccacag	ccccagagaa	5040
gaaagggtta	tttcagacat	tctctctaat	atgcattgaa	ccattctgaa	ttttaccagg	5100
ttctcgtctg	agcaggatct	ctattgagaa	aaagttaggg	tcagtaaggg	ggaagggtct	5160
gtccacagat	gaagtccaat	tcgattaaag	gggataaggg	aatacatgtg	ctcttagctt	5220
gaccaggtag	ggcaaaagg	gaagcatata	tgaaggcagc	ttcagaaaaa	ctcaagctgag	5280
cactgacttc	agactggaat	taggaatcca	gcctctgccac	tttattctac	tcagcaataa	5340
tttactgagc	aaattctatg	ggctagacag	tggattgggt	tcacaagata	caatgagtgt	5400
gacatgtgtg	ttgtctatgg	atttggggat	atatgtaggt	atagggatat	cttcaagggt	5460
aaatcaaggg	ttctaatgag	gccagccatg	gtggctcaca	cctgtaatcc	cagcaatttg	5520
ggagaccgag	gcgggtggat	cactcgaggt	agactagctc	caggactagc	gaccaaacat	5580
gtgaaccccc	gcctctacca	aaaatacaaa	aattagttgg	gcgttatggc	aggctcctgt	5640
aatccccagt	tctcgggagt	ctgaggcagg	agaattgtct	gaactgtggt	ggcagagggt	5700
gcagttagcc	gagattgttg	ccactgcatt	ccagcctggg	tgacagagcg	agactttgtg	5760
tcataaaaaa	aaaaaaaaga	aaagaaaaga	aaaaggagct	ctaatgagat	aaaaatgaaa	5820
aagcctggca	tgtagtggca	acttatgaaa	aattgttaatt	aaaaaaaac	atttctgcag	5880
agaagaaact	ggatctacct	ggtttttctg	aagcctaatt	ctgctcgccc	cagtgagtgc	5940
tgtttctgag	gcattccgtg	tgttttgagc	tgtggattgt	gaaggttaga	gtgggaggga	6000
tttttagaggt	taggtctgcc	cctcttctgt	tagaggacat	ggatccctgg	cttgaggagg	6060
ttctggtttt	tgtatcaagc	ctcacaaggg	gtggcaccac	ctcactctta	ggaactccgc	6120
tagaagggaag	gccagctctg	cctaattcgg	ttggggagat	gggggtccct	ttatgctagc	6180
agaaatgtct	cgaaggagca	tgatgggtgc	agctttgttc	atgaaggcca	gtgtgcacaca	6240
ggagaccggg	cagctctctc	agcagctccct	gtggccaact	ttccttaagt	cttgaggagt	6300
cttttttttg	cacaatctca	gtcactgca	acctccgctc	cccaggttca	agcgattctc	6360
ctgcctcagt	ctccacagtg	gctgagacta	caggcatgca	ccaccaagcc	cagctcaatt	6420
tatatatttt	agttagagat	gggttcacca	tgttgccacg	gatggtctcg	atctcttgac	6480
ctcatattcc	acctgcctcg	gcctcccaaa	gtgctgggat	tacaggtctg	agccactgag	6540
ctgcggccag	gagttcttaag	ctgagatcac	agcattgcac	tcacagctgg	gcataaagag	6600
caaaaactca	tctcaaaaaa	aaaaaaaaga	tagacacaag	actggctctg	tgtctttttt	6660
ggggacaggg	tctcactcta	tcacccaggg	tggagtgcag	tggtgcacac	acagctcact	6720
gcagcctcga	tttccaggcc	tcaagtgaac	ctccactctc	agcctcctga	gtagctggga	6780
ctacaggtgt	gtgcagctga	gcctggctaa	tttttaaaaa	ttttttgtag	agatgaggtc	6840
tcactatatt	ggctgggggg	cctcaaaact	ctgggctcag	cagtcctccc	acctcagcct	6900
cccaaaaggc	tgggattata	tgtctgctct	ttttaagggt	ctttagaggga	caaacatttc	6960
acctactcct	tgtcaagcca	gtggaccggt	gggtccagac	atacggctaa	agtcaaggag	7020
tgtgtctctt	tggagaataa	ctttcaatca	ggaatttcaa	tcagaaattc	aatcatgtgg	7080
agagagactt	atctcaaaaa	tgtggtgggt	cgtgggatgc	tctgttttat	tagttccttg	7140
acagtatgta	tgtgtgtgag	tgtgtgtgtg	tgcgcgcga	cactcatttg	gatgggtgtg	7200
tatgttgttg	ggggggtggt	gcgtacgtat	gtggatgtgt	ggatgtgggt	tgtgggtgtg	7260
cgcgctcata	ggtggagggt	tgtgtatggg	tgcgggtatg	tgtgtgtgtt	ggggactggg	7320
attttacag	ctctccaggg	gctgagtga	ggctttcggg	caagctcctc	gggagctagg	7380
caaaagctgag	tgtattctct	gttatgccat	ttattattgg	gttgcaacctg	gtgaaactgc	7440
caatattctca	cactttgact	tttatttatt	ttatttttta	ttttttttga	gacgagagtt	7500
ccacttgttc	acctaggctg	gagtgcaagt	gcggatctc	agctcaactg	aacctctgcc	7560
tcattgattc	aagtgaattc	cctgcctcag	cctcccaagt	agctggaatt	acaggtgccc	7620
gctaccacgc	ctgactaatt	tttgtatttt	tggtagagac	gggattttcac	catgttgttc	7680
aggctggctg	gaaactcctg	acatcaggta	atccacccac	ctcagcctcc	caagtgctgt	7740
ggattacagg	catgagccac	tgcgcocggc	cttttttgac	ttttaaaaat	gggagtttga	7800
ataatactaa	tcaggtggtt	gaattagcta	gcactgattc	ctgcacaaat	ctcaggtttc	7860
ctcacagctt	agagtcaaaa	gcagggctat	gggaattcca	actaaaaata	attttgaaaa	7920
tgccttatga	aattatcact	ccaaagaaat	gcgcgcagtg	tcagttgtct	ctgtctcttc	7980
atctcacagt	atcacatttc	gcggattagg	aagctgagtc	tgagangctc	cgtgtgtgtc	8040
tttttcggag	gcaccgtgat	gtgatggga	gctcactcgt	taggaagtca	gaacagagtc	8100
tctgagggat	catttctcta	atctgtcagt	ttcctcatct	ctgaagttgg	gctcatttcc	8160
ttcctctcag	gagttattgt	aaagatgaag	ataaaataag	tgtaaaaact	agcatgggaa	8220
ctggcttcta	taaggttcta	ataagtgcat	ttcctactct	ttccctcagc	cttcccaatt	8280

gtaaaagcaa	ggcagggggt	aggtgatttc	tggggctcct	tttggctctg	acatttgagg	8340
attttgtatc	cttttttttt	tcagagtctt	gctctgtcac	ccaggttgga	gtgcagctca	8400
atgcaaatc	cgctccccc	gctcaagcaa	ttcttatgtc	tcagcctcct	gagtaacctg	8460
gattacagcg	aggcaccacc	accocagct	aattttttgt	attttcaagt	gagacggggt	8520
tttgccatat	tggccagggt	gggtctgaac	tcctgacttc	atgtgaccca	cccatctcgt	8580
ctctcccaag	tgtgagatg	acaggttgga	gctacccgtg	ctggcccaat	tttgtgtgct	8640
taattcgctt	ttctgttgga	agagttggca	ccaggttttg	tgtatctctt	ccccacaag	8700
gctctgcctc	ctgcccagtc	cagaggggac	cctgtccttg	caatttcacag	gattctcgtc	8760
ttgcaactga	aattccagta	ggtcaaaagt	aaattttoca	tacacttttaa	catgaagata	8820
aatgatcaca	gtatggccct	ttagatctct	gagaacatca	cggtcatccc	ctgtgataat	8880
tttaaaagca	gatgaatcca	tgccctgtcg	aggtttgcca	ggaaagccag	tgctgggatt	8940
acagtggag	tctttttatg	ctactttttt	cttgatcccc	tcaccccatg	gggtggcata	9000
ttgaaaggca	ggatgtgtga	ccacgatact	ttctctccct	tggaactatg	ctaagagtct	9060
gttattgggt	tctgaagatc	agagtttaat	ttccgaactc	tctctgtgta	gctctgggat	9120
cttggaagag	cacttaacct	ttctgaagtc	ccctttccct	atctctaaaa	tgcatacact	9180
catcactaac	atttactgag	cactgacatg	tgcagacac	catcttaagc	attttacaca	9240
gactcacaca	tttgatcttc	caacaaacag	aaactatgaa	cgcatctacg	gtcagaacaa	9300
atgattttgt	octaagcacc	aagaccgtag	agccctgtct	ccctattcta	ccctactcgt	9360
tctctcaaaa	tgatttgtgag	aatcgatga	gacactagg	gagaaaagg	ttttataaat	9420
agcattttta	aaatttttta	aagtcacaaa	aaattttta	tttaatacag	ataaaataga	9480
tccttttgtt	ttataaaaa	taacaaaatt	tgttatacaa	caactatgtg	attttataat	9540
tttgcttttt	tgtatgtctg	caggaaagaa	acattaaaga	atcttaaat	gattatctgt	9600
aatcagaagg	tctgctctga	cttttttatg	ctcctaactg	acagctgatc	atacactctc	9660
attttttttt	atgacacttc	aagggtgcgc	ttagcttcat	caactcctctg	ttgcacaaag	9720
ctttgtgacc	aaaaaacatt	aagcagattc	ctgagtcact	aaatgacaca	taaccagagt	9780
tgagacttag	gaacttttag	tgccatgcta	agccacaggg	gacacacaaa	atagcatttt	9840
acaaaggcaa	agaattttga	caacttgagat	ttagcttggt	gatccttgta	aaagtttttt	9900
ttttaggcat	aatttaggtt	tagatcatag	tactcactat	tacttagtaa	taattttttt	9960
ctgatagaaa	tacagtgtaa	caggccgggg	gcagtggctc	atgctgtgaa	tcgccagcat	10020
ttgggaggcc	gaggcggggc	gatcacttga	ggctcaggat	ttgagaccag	ccctggccac	10080
atggtgaaat	cccatctcta	ctaaaaatc	aaaaaattag	ccaggtgtgg	tcgtgtgatt	10140
ctgtgatccc	agctactgtg	gagggtgagg	caggagcatc	agttgaacct	agggaggcga	10200
ggttcagctg	agcccaagat	gtgccattgc	actccagcct	gggacacaaa	cgcgactctc	10260
acttcagaaa	caaaaaaaaa	aagagagaga	gagaaaagaa	gaggaaggag	aggaaggaaa	10320
ggaaagaagg	aagggaagg	ggaaagaagg	aagggaagg	ggaaaggaaa	aagggaagg	10380
ggaaagaagg	aagggaagg	agaagggaag	aagggaagg	ggaaaggtaa	caagcaaatg	10440
gtatacaatg	caatatctaa	aaaaataggt	atttttatat	gtttgtcggt	ttatatatat	10500
gaccccccac	tttagagatg	ggaaaactgag	agattaagg	aacgatccct	gagagactct	10560
gtctcgactt	ccaaactcgt	gagctttcca	tcgcatacag	gtgcctccga	aagcatgaca	10620
cgagagctct	cagacttagc	ttcctaactc	gctaaacggg	attatgtgag	atgaggaaca	10680
tggaaacgct	gacatgggtg	aggggtccct	ggagtatcat	tttcatgtgg	cattttcaaa	10740
acttatttta	cctaactctc	ccaaagccct	gcttttgact	ctaattgtgc	tcctgagact	10800
tggaagagcg	aagatgctag	gcacagagca	agactccatc	tcagataaaa	taaatagaat	10860
aaaaaaaaaa	gaacacaaat	aaatttgaaa	atttttttga	aaattaggca	cgtttgcaat	10920
gaccttctaa	tgttattaat	tgtctggttc	ccaccacaga	ttagtttgga	atgcaacttt	10980
cttttacaat	cagagtccgt	tcttggtctt	ggaaactctt	gaggtcctct	tgcatctccc	11040
actcttgat	ttttggcaac	tctacccctg	gccactgtca	tggaacccag	gctgatcgca	11100
cotatttagtg	gagaaatmtg	tcataatac	tgaagtgttg	ggacaaacag	tgctccctta	11160
gggttaggaga	aagagatctt	tatttttaac	aaagggggag	gagccagaaa	actccagaga	11220
ccccagagct	tgccctctct	ccaaagtttg	gggtaaaccc	cccgtaacc	ttatctctct	11280
gggctttcac	atattctgga	ttctctccct	ctgtttccca	gcagaaaagg	atggagcctc	11340
acagattctt	cccatctctg	gagaaaaaca	tgcatggagc	tcaaagttct	tctcaggagt	11400
tttattgcca	aagcacaatc	aagaaagggt	agcagtggag	agcagtggag	aagttaaaag	11460
atgcataaaa	tctgttaaagt	ctcagacaaa	gaattctcct	aaaatgcaaa	aggggctctg	11520
ctgctctccc	cttggtctctc	catgtagctc	tcttatctct	tgcttatctg	agactagtca	11580
aaactaagct	gtttctctat	ttatttccag	aagctattga	gaacactctc	ctgaattctt	11640
caaaatcagt	agagggcgac	aaatgtacat	ataaatgatg	tgagtgggtc	ttaaataaag	11700
actcatgaca	ctaaaagggt	cagcacctgc	gtctgattgc	acctgtttct	gtgctgtctt	11760
ctgtctctct	tctctctctc	tgccatttca	ttatcaatgg	ttactttact	tataagatac	11820
tattagaacc	tgtatattga	taaatgatgc	atcagatcta	tagtgagaga	aaaaattaat	11880
gcaattaaag	gtgtgtgaac	agctagctct	caagtgggga	gaattcattt	gagtaacctta	11940
ggctcagact	tacatcaaaa	caaaaaatca	gagctacatt	aaaaagtgaa	tttttaacta	12000
tatcaaaaca	tagaaaaaaa	cagaagaaaa	ttgaatactt	actaaatctt	agcatgaata	12060

agaactgttt	aacacttaga	ggcaaggact	ggcggtggtg	gctcatgctt	ttaatcccgag	12120
gactctggga	ggccaaaggcg	ggcggtatcac	ctgaggtcac	gagtttgaga	cgacgcctggc	12180
caacactggcg	aaaccccgctc	tctactaaaa	aatgcaaaaa	ttagctgcgt	gtggctgggtgc	12240
aaactgtgta	ccatgctcac	ttgggaggct	aaggcatgag	aatcgctgat	acctggggagg	12300
tggaggctgt	agtgagccga	gatttgtcca	ctgcactaca	gcctgggtga	cagtgtgaaa	12360
tccctctctc	caaaaaaaaa	aaaaaaaaaa	agcaaaactag	agcagtgagg	taccattatt	12420
tcccttgctc	actaaactga	caacacacaa	atgtttttta	taatacccaa	gagctgtgag	12480
ggtagttaa	gtatgcocct	ttatacacac	actaatgatg	tactactggt	tggcagatata	12540
acatatgctg	ccatgtgggg	atatgtatca	ggagagactaa	aaatgtgcac	acocctttggg	12600
ccagtaaatc	actttcggaa	atctgtcata	acagaataat	aatcttgggg	agaagtcatac	12660
gcctcaaggat	atttcaaaaa	ttatttaaaa	atcaaatgat	aatctcttac	agaatatataa	12720
ataatatctt	aaaatgaaaa	tatgctaaaa	gtttgatgaa	atataaatgg	tcaaatatac	12780
attgattata	tccacttaact	agactagcac	tcaactcgag	acgttataaaa	tagtctattat	12840
aaaaactcaga	aaaatgcaaa	gacaaaaata	aggaataaag	ttttacataa	agatgatctc	12900
cactcatgttt	aaaaatpaac	agagacattc	ttggagttga	gtattgtttt	cttttctcgtc	12960
atgtccaaag	aaactataaa	ctatttatttt	taatgaacta	tatatgtata	atatacatat	13020
agtttatatg	tatatcaaaa	atttatctca	tatatatgat	aaagatgaaa	gatgatgttgg	13080
atgtgccagc	tgaagtgggt	agtatagaaa	ccaggttaat	ggggcatagg	agtggtgatc	13140
cagataccag	gcccattgtt	ttgggggtgag	attgccaatc	acogtctctc	ttccatccct	13200
ccagagggag	taggtttgtg	ttcaacaaac	cttcagttgt	ctcgaagaca	aacctaattc	13260
tggagacttc	tataaatcta	gaagagacaa	gcaaaactgat	gaaaaatagt	gaatttttaa	13320
ggtaaaataa	agtacatact	ctcacactttg	tttagaatca	gattctttggg	atbaaccaca	13380
ttaacccaca	gagggctgtga	gtgatgcctc	taatccaggga	tcctaggacc	tatttctctc	13440
tgtgagatgc	ttcttcccaa	ctccttggtg	agagtgaggaa	gactaagaac	tcagcaaatct	13500
ggatggggag	ccctaagatc	cttcagagac	ggagggcagaa	tcctgagaggg	gataaaagtc	13560
cctatacctg	tattggggcc	ttttctggga	gggggatatc	aaagaatgat	ttttgagacag	13620
ggagggcttt	gtactacctg	gccacttgag	cttttgcata	gggctccaga	atcacatattt	13680
caaaatacat	ccccctccct	ctctctctcc	ctcttccact	cttctctttt	atcttctctt	13740
cttctcttct	ttctctctct	ctctctcttc	ctctgctctc	tcattgatctt	ttctctctct	13800
tataaaagtc	cttatttagt	ccctactctg	ctatttagtg	gttagtctct	gtctccctgg	13860
acttgtctgt	taatggagaa	atgggtgagc	aaaacagaaa	tacacagcaga	gtgcaataat	13920
agagctaagc	caggtgtata	aatccattct	cacactgtct	taaaaaacta	ctgggtaatt	13980
tataaagaaa	agaggttttaa	ttgactcaca	gttccacagg	ctgtacagga	agcactgaagt	14040
gggagccctc	agaaaactta	caatcatggt	gggaagaaaga	gcgaaggggga	gcagagcaca	14100
tcacacagca	gcagagagaga	gagagagaaa	gagagagaga	gagaatatag	gggaagtgtct	14160
acacactctc	aaccagattct	tgtgagaatt	caocactact	catgagaatac	gcaagggata	14220
agctgcctc	catgatctag	tcactcctca	ccaggccctc	tctccaacac	agctgcagct	14280
gctattttgg	tggggacaca	gaccocaaac	atattaccag	ggcactggag	aaacacagag	14340
gggaagaagac	cagccaaagg	gtgagatgga	gaacaaaggag	gactcttctga	aacagatgac	14400
atccocaaact	ggctctgaaa	gctgaataga	gattagacag	gggagggagg	gcagctaaaag	14460
atggctcagg	caaaaagagg	gccaggggat	atgttcatgg	gatgatgtgt	ctctctgtgt	14520
ctgcttaaca	caaggttgagt	ctctccctcc	ctctctctct	cttttctctct	gtgtgtgtgt	14580
gtgtgtgtgc	atgtgtgcga	atgtaatatata	cccaatagtct	aaacatgtgc	ccagagagag	14640
gggtagagga	agaaagagaa	tgagagagta	agaaggagga	atagacacag	aaaatgagag	14700
agaagggggg	aaagaaaaag	aaagaaaggag	ccagaggaga	gaagctgggt	agcattgaaat	14760
ggagcaatct	gtgtcatcgt	acttgggaaa	cccaaggatg	gattcttggc	aagtgcagctc	14820
ttggagctct	ccctgtgtct	ggctctgtgc	tcagacatgg	gaaaattaga	ggaggtgtcat	14880
ctgtgcaact	actgaattca	taactttggt	gaggaaagga	gactacacaa	aggggaataat	14940
ctcgaagtatt	acagatttca	gggcagaaag	agatcaaggt	gggctgcaat	attcagaaaa	15000
gtctctctgg	aaaagtgtgaa	tacttagaaa	gcagctccta	gaagtagact	ctgtgtagat	15060
ggaaggagtc	ctttgttaggt	cccaactggg	tgtgtgtgtg	gggtctgtct	ctccaaggcy	15120
gcagctgcac	atgtggattc	cagggtctcag	gatgctgtgt	ctgggagctg	ttctactgct	15180
attagctctg	ccgggkcatg	accaggaaac	caagactcaa	ggggcccgag	tcctgctctc	15240
ctcggcccaag	ggggcctgca	cagggttggt	ggcgggcatc	ccaggggcatc	cgggccataa	15300
tggggcccca	gggocgtgat	gcagagatgg	gagaagtggt	gagaagtggt	agaaaagaga	15360
tccaggtlaag	aatgtttctg	gocctcttca	tcacagacot	ccacacatga	tataaactat	15420
atgaagkcat	ctattattaa	ctaaaggcta	gaacagggga	caaaagcaaa	ctttttctgt	15480
ttaacctata	gcacactgag	gtgattttgg	gttgtctctc	caaggatgag	tgtgatgtgt	15540
gcctctataa	ccaaagactt	ggctttgctg	catctgcagc	ctcttttcca	tcocctctcc	15600
catcttccac	ctcatcccta	ttccagatc	atctcatctc	tgattctctc	ttctgtctgc	15660
ttaactttca	tttccaccaa	tggcaltcaa	ccacactttc	tgcacacccc	ctgaaggagct	15720
cagctctgco	tttgggaaac	tcttgatcta	ggtaagatgt	ctaagtgtca	aggtctctgt	15780
ggtggttacc	acagaaaggt	ctactctaaa	aatgtcaaac	tgaattgtgaa	caagtattca	15840

aagtagtgag	catagagaaa	atrtactcac	cgtggacctg	atgaagaatg	aagcgttcaa	15900
ggagggaggca	gagccttcagc	taggccttga	atgatgggta	ggcagaatga	aggagggagag	15960
acatcctgata	tggagggggt	agaattgcga	aaccagggtt	gatgggtgca	gcacataaag	16020
ggctggcagg	gtggagggtc	tatgatagag	acctatagga	gataaagata	gagttgaaat	16080
tatgggagcc	tcocatgtctg	tgggagatat	agaaggagga	ggtaaacacct	ctctcctttt	16140
gggagctctt	attgtgtttc	tgatctataa	gtcaagaagg	ttgtgagtgg	gagccacagag	16200
gagggtaatt	taggctgttaa	ccaacctagg	caggagttct	gtctcttgta	gtccatgagg	16260
ctctctcatt	ccttaggtct	tatttggtct	aaggaggaca	tcggtgaaac	cggagtaccc	16320
ggggctgaag	gtcccggagg	ctttccggga	atccaaaggca	ggaaaggaga	acctggagaa	16380
gggtccatag	tataccgctc	agcatttcagt	gtgggagtgg	agacttacgt	tactatcccc	16440
aacatgcoca	ttcgctttac	caagatcttc	tacaatcagc	aaacccaata	tatgtggtcc	16500
actgggtaaa	tcocattgca	cattcctggg	ctgtactact	ttgcctacca	catcacagtc	16560
tatatgaag	atgtgaaggt	cagcctcttc	aagaaggaca	aggtctatgt	cttcacctat	16620
gatcagtagc	agggaaaaat	tgtggaccag	gcctccggct	ctgtgctcct	gcactctggg	16680
gtggggcagc	aagtcctggc	ccagggtgat	ggggaaaggag	agcgttaatg	actctatgct	16740
gataatgaca	atgactccac	cttcacaggc	ttttctctct	acctgacac	caactgatca	16800
ccaactaact	agagcctcct	ccaggccaaa	cagcccccga	gtcaattaaa	ggccttcagt	16860
acggttagga	agtttgattg	tatttagttg	gagggcctta	gtatatttca	attcattttac	16920
tacttctatt	attcattcct	tcatcaagta	actttaaaaa	aatcatatgc	tatgtcccca	16980
gtcctgggga	gcttccacaa	catgaccaga	taactgacta	gaaagaagta	gttgacagtg	17040
ctatttttgt	ccaactgtct	ctcctgatgc	tcatatcaat	cctataaggc	acagggaaca	17100
agcattctcc	tggtttttaca	gattgtatcc	tgaggctgag	agaggttaag	gaatgtctaa	17160
gtgcacacaa	gtattaaagt	acagtgctag	aaatcaaaa	cagagctgtg	gactttgttc	17220
actagactgt	gcccttttat	agaggtacat	gtttctcttg	gagtgttggt	aggtgtctgt	17280
ttcccaacct	acctgagagc	catggaattt	gccttcctca	tgatattaaa	ctcccccaca	17340
cgagagcttc	ctcagagaaa	gtggttctat	gatgaagtc	tgtcttgaaa	ggaactactac	17400
tcaatggccc	ctcactactc	ctactctctc	ttactatgt	cccttctcat	gccttccctc	17460
ccaacgggga	aagccaactc	catctctaag	tgtggaactc	atccctgttc	ctcaaggcca	17520
actggccagg	agcttctctg	atgtgatata	caactttttt	tttttttgag	atggagctctc	17580
cctctgtcac	ccaggctcga	gtacagtgac	acgactcggg	ctcactggag	gccttctctc	17640
ctggggtcaa	gcaattattg	tgccctcagc	tcocgagtag	ctgagacttc	aggtgcattc	17700
caccacacat	ggctcaattt	tgtattttta	gtagaataag	ggtttcgtca	tgttgccocag	17760
ctggtgtctg	aactcctggc	ctaggtgatc	caaccgcctc	gacctcccaa	agtgctggga	17820
ttacaggcat	gagccacact	gccagtcga	tatctcaact	tttattttgc	ctcaggttag	17880
agtcctgggt	gtgaggaaaca	ctcccccaca	ggctagaggc	aactgcocag	gaaggactgt	17940
gcttcctgca	ctctcaaatc	ccttgacagt	ctcctgataa	tgctcatatg	agaccaaatc	18000
cttgaatccc	gtatctaccc	agaattaaat	ccattccagt	ctgtgcattg	aatcagtttt	18060
actccacaga	acattttctc	tttaggaaat	ccctggtttt	aagtatcaat	ctctgttcag	18120
ctggacaata	tgaattcttt	ccaactgaag	tagggatgac	tgtgtatttc	agaaacagtc	18180
cagaattttt	catcaagaag	gtagccttgag	cctgaaatgc	aaaacccatg	gagaaattct	18240
gaagccactg	ttctccttag	taccaacagg	gtcaggggaag	actgggcctc	cttgattttat	18300
tattgtctct	taagaaatag	aggttgaggt	agttgtatgt	ggtaaacatt	ctctcagggg	18360
acaaataact	cagtgatttct	cttcaagagt	tttagcaaaa	acagagtata	tgaactcttc	18420
tatcaatata	taaattttaa	aaactatctt	tttgcttaca	gttttaaatc	ctgaacaatt	18480
ctctcttaca	tgtgtattgc	taactattaa	ggattatttt	tttccacata	taagagcttg	18540
tccttttgtt	gtgtgttttg	tttttaagat	ggagtgttcc	ttgtgtgcca	gcctagatgt	18600
cagtggtcgt	atctcggtct	actgcaacct	ttgcctccca	ggttcaaggc	attctctctg	18660
ctcagctctc	cgagttagct	ggacacacgg	tgcctaccac	catgcocagg	taatttttgt	18720
attttttagt	aagacagggt	ttccacattat	tggccaggct	ggctctgaac	ctcctgacct	18780
gtgatctgcc	caactccatt	ttttgttgta	ttttttgaga	aagatagata	tggagtttag	18840
agagggatga	agaggtgaga	gtaaagcctg	tgttagtcag	aactctgtgt	tgtgaattgc	18900
atgcacacaa	gaaaaccocaa	aatattatgc	aaactactgt	aagcagaagaa	ataaaggaaa	18960
aaatggaaac	atttattctc	tgtcataata	gaaattacca	gagttgtctc	gtcttagatg	19020
aaggtttgaa	ccaaagctca	aaacaatcaa	gaccttttc	tgtatgtcct	ctctgtctgc	19080
cttcocgagt	gtagggttta	ccctcagggt	ctacacagta	tagtattcag	gtttccctcc	19140
cgatatacaa	aagactctgg	cctgccacgc	tctcgtatac	ccaagccaca	ccatctgggt	19200
aaatggacat	catgtttctc	gggtgatccg	aaagaggaga	gagggaagtc	ttctccocag	19260
atgccccagc	aagtgtaacc	tgtcatctca	ttgctctggc	ctgatttgctg	gcctgtttct	19320
gaccacacac	tgagtccagg	ggatgaataa	ttcatattga	tttaatttga	ctcttaagta	19380
gggttatgta	gaggtatttt	ccctaaagca	aaattgggac	actgttatca	gaaataggag	19440
agtggtatgt	agatgcacaa	taataacctg	ccacaacaaa	ctcttaatgt	tgtgttttag	19500
ttctcatgag	tttccagag	agacatagct	ggaaaattcc	tattgatatt	ctctaaaaat	19560
tcaacaagta	gctaaagctc	ggctatgctc	acagctcac	atctggttgg	gggtgggtcc	19620

ttacagaaca	cgctttcaca	gttaccctaa	actctctggg	gcagggttat	tcctttgttg	19680
aaccagaggc	acagagagag	tcaactgagg	ccaaaagagg	cctgagagaa	actgagggtca	19740
agatttcagg	attaatgggc	ctgtgatgct	tgaagtaca	attgtggatt	tgctcaattc	19800
tccttagtgc	tgctcagctt	tgcttcata	attttagcgc	tctattatta	gatatataca	19860
tgcttagtat	tatgtcctat	tggtgcattt	actctcttat	cattatgtaa	tgctctctct	19920
tatctgtgat	aattttctgt	gtctggaagt	ctactttgtc	taaaaataac	atacgcactc	19980
aacttccttt	tctttctctc	ttcctttctt	tcttctctcc	ttctctctct	tctctctctc	20040
tttctctctc	tccttctctc	ttttctctct	ctctctctct	ctctctctct	ttttgacaga	20100
ctctctctct	gtggcctcgg	ctggagttca	gtgggtgat	cttggctcac	tgctacctct	20160
accatgagca	attctcctgc	ctcagcctcc	caagtagctg	gaactacagg	ctcatgccac	20220
tgcgcccagc	taatttttgt	attttctgta	gagacggggt	ttcacccat	tcgtcagggt	20280
gggttcaaac	tctgaacttt	gtgatccacc	cgctctggcc	tcccaaatgt	ctgggattac	20340
aggcatgagc	catcacacct	ggtcaacttt	cttttgatta	gtgttttgt	ggtatatctt	20400
tttccatcat	gttactttaa	atatacttat	attattgtat	ttaaaatgtg	ttctctacag	20460
actgcattga	gttgggtata	atttttatcc	agtctaaaaa	tatctgtctt	ttaattgggtg	20520
tttagacaat	ttatatttaa	taaaattggt	gaattttaaga	tggtgactgt	ttttatttgt	20580
ttgctgttca	ccacttctgt	tttattctct	ttccagaatt	cttttggtat	gtttaaatat	20640
ttcataatat	tttatcttaa	tttatattat	gggtatttgc	ctatatctct	ttgtggtatt	20700
tttttagtgt	tgcttgaggg	attacaattg	acttaacttt	tccagtggtg	cataaagtta	20760
atattttggc	acttgcaagta	aaccgtagaa	ggcttataat	catattagta	cctctatcca	20820
ctttctttta	tggttgtagtt	gtcatatata	ttacatctat	atacactgaa	acattatagg	20880
caatgttatg	atttttgcat	tcgtcagtc	tatatatat	ttaaagaatt	taagaggaga	20940
aaaatacata	ttcagatat	catcat				20966

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 May 2003 (30.05.2003)

PCT

(10) International Publication Number
WO 2003/044057 A3

(51) International Patent Classification⁷: **A61K 38/22**,
A61P 25/28

(21) International Application Number:
PCT/IB2002/004924

(22) International Filing Date: 29 October 2002 (29.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/332,119 21 November 2001 (21.11.2001) US

(71) Applicant (for all designated States except US): GENSET
S.A. [FR/FR]; Route Nationale 7, F-91000 Evry (FR).

(72) Inventor; and

(75) Inventor/Applicant (for US only): LUCAS, John
[US/US]; 339 The Valley Road, Concord, MA 01742
(US).

(74) Common Representative: GENSET S.A.; c/o De Luca,
Giampiero, Serono International S.A., Intellectual Prop-
erty, 12, Chemin des Aulx, 1228 Plan-les-Ouates (CH).

(81) Designated States (national): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
10 June 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF OBG3 FOR PROMOTING CENTRAL NERVOUS SYSTEM REMYELINATION

(57) Abstract: The present invention relates to the field of central nervous system (CNS) research. Demyelination of neuronal axons within the CNS underlies the pathogenesis of degenerative diseases of the neuromuscular system, such as multiple sclerosis and hereditary leukodystrophies. Therefore, treatments aimed towards accelerating the repair of myelin sheaths offer a potential therapeutic to ameliorate the symptoms of multiple sclerosis and leukodystrophies. A compound, globular OBG3, has been identified that has immunosuppressive properties. This compound should be effective for accelerating the rate of remyelination and treating multiple sclerosis and leukodystrophies.

WO 2003/044057 A3

INTERNATIONAL SEARCH REPORT

PCT/IB 02/04924

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K38/22 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

MEDLINE, EPO-Internal, WPI Data, EMBL, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>ALLAMARGOT C ET AL: "A single intracerebral microinjection of platelet-derived growth factor (PDGF) accelerates the rate of remyelination in vivo." BRAIN RESEARCH. NETHERLANDS 9 NOV 2001, vol. 918, no. 1-2, 9 November 2001 (2001-11-09), pages 28-39, XP002236251 ISSN: 0006-8993 the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/-</p>	1-11

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

3 April 2003

Date of mailing of the international search report

07/07/2003

Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Lechner, O

INTERNATIONAL SEARCH REPORT

PCT/IB 02/04924

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	YE P ET AL: "INSULIN-LIKE GROWTH FACTOR-I INFLUENCES THE INITIATION OF MYELINATION: STUDIES OF THE ANTERIOR COMMISSURE OF TRANSGENIC MICE" NEUROSCIENCE LETTERS, LIMERICK, IE, vol. 201, no. 3, 1995, pages 235-238, XP001149310 ISSN: 0304-3940 the whole document ---	1-11
A	HALLEUX C M ET AL: "Secretion of adiponectin and regulation of apM1 gene expression in human visceral adipose tissue." BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS. UNITED STATES 16 NOV 2001, vol. 288, no. 5, 16 November 2001 (2001-11-16), pages 1102-1107, XP002236252 ISSN: 0006-291X the whole document -----	1-11

FURTHER INFORMATION CONTINUED FROM PCT/ISA 210

Continuation of Box I.2

Claims Nos.: 1-4, 7-11 (only partially)

Claims 1-4 and 7-11 (as far as relating to in vivo methods) are not acceptable under Article 6, PCT. The therapeutic application is functionally defined by a result to be achieved and pathophysiological mechanisms, respectively, i.e. regeneration or myelin sheaths of axons, increasing the rate of cellular division/migration of progenitor cells, inducing oligodendrocyte progenitor cells to differentiate into mature dendrocytic cells at sites of demyelination in axons and diseases in which demyelination of nerve fibres occurs.

However, this does not allow any practical application in the form of a defined, real treatment of a pathological condition (disease). Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely the specific diseases listed in present claims 5 and 6.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-9, 11 (only as far as relating to in vivo methods) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-4, 7-11 (only partially)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.